arXiv:1210.0031v1 [math.OC] 28 Sep 2012

OPTIMAL CONTROL OF A FREE BOUNDARY PROBLEM: ANALYSIS WITH SECOND ORDER SUFFICIENT CONDITIONS

HARBIR ANTIL*, RICARDO H. NOCHETTO†, AND PATRICK SODRÉ‡

Abstract. We consider a PDE-constrained optimization problem governed by a free boundary problem. The state system is based on coupling the Laplace equation in the bulk with a Young-Laplace equation on the free boundary to account for surface tension, as proposed by P. Saavedra and L. R. Scott [16]. This amounts to solving a second order system both in the bulk and on the interface. Our analysis hinges on a convex constraint on the control such that the state constraints are always satisfied. Using only first order regularity we show that the control to state operator is twice Fréchet differentiable. We improve slightly the regularity of the state variables and exploit this to show existence of a control together with second order sufficient optimality conditions.

Key words. sharp interface model, free boundary, curvature, surface tension, pde constrained optimization, boundary control, lagrangian, control-to-state map, existence of control.

AMS subject classifications. 49J20, 35Q93, 35Q35, 35R35.

1. Introduction. Free boundary problems (FBPs) are challenging due to their highly nonlinear nature. Besides the state variables, the domain is also an unknown. FBP find a wide range of applications from phase separation (Stefan problem, Cahn-Hilliard), shape optimization (minimal surface area), optimal control problems with state constraints, fluid dynamics (flow in porous media), crystal growth, biomembranes, electrowetting on dielectric, to finance. For many of these problems there is a close interplay between the surface tension and the curvature of the interface [21, 22].



FIG. 1.1. *Ω_γ denotes a physical domain with boundary $\partial\Omega_\gamma = \Sigma \cup \Gamma_\gamma$. Here Σ includes the lateral and the bottom boundary and is assumed to be fixed. Furthermore, the top boundary Γ_γ (dotted line) is "free" and is assumed to be a graph of the form $(x_1, 1+\gamma(x_1))$, where $\gamma \in \mathring{W}^1_\infty(0,1)$ denotes a parametrization. Γ_γ is further mapped to a fixed boundary $\Gamma = (0,1)$ and in turn the physical domain Ω_γ is mapped to a reference domain $\Omega = (0,1)^2$, where all computations are carried out.*

Of particular interest to us is the control of a model FBP previously studied by P. Saavedra and L. R. Scott in [16] and formulated in graph form; see Figure 1 where the free boundary Γ_γ is the dotted line. The state equations (1.2b) involve a Laplace

---

*Department of Mathematical Sciences. George Mason University, Fairfax, VA 22030, USA (`hantil@gmu.edu`).

†Department of Mathematics and Institute for Physical Science and Technology, University of Maryland College Park, MD 20742, USA (`rhn@math.umd.edu`).

‡Department of Mathematics, University of Maryland College Park, MD 20742, USA (`sodre@math.umd.edu`).

equation in the bulk and a Young-Laplace equation on the free boundary to account for surface tension. This amounts to solving a second order system both in the bulk and on the interface. Below we give a detailed description of the problem.

Let $\gamma \in \mathring{W}^1_\infty(0,1)$ denote a parameterization of the top boundary (see Figure 1) of the physical domain $\Omega_\gamma \subset \Omega^* \subset \mathbb{R}^2$ with boundary $\partial\Omega_\gamma := \Gamma_\gamma \cup \Sigma$, defined as

$$
\begin{aligned}
\Omega^* &= (0,1) \times (0,2), \\
\Omega_\gamma &= \left\{(x_1,x_2) \,:\, 0 < x_1 < 1,\ 0 < x_2 < 1 + \gamma(x_1)\right\}, \\
\Gamma_\gamma &= \left\{(x_1,x_2) \,:\, 0 < x_1 < 1,\ x_2 = 1 + \gamma(x_1)\right\}, \\
\Sigma &= \partial\Omega_\gamma \setminus \Gamma_\gamma, \\
\Gamma &= \left\{(x_1,x_2) \,:\, 0 < x_1 < 1,\ x_2 = 1\right\}.
\end{aligned}
$$

Here, Ω^* and Σ are fixed while Ω_γ and Γ_γ deform according to γ.

We want to find an optimal control $u \in \mathcal{U}_{ad} \subset L^2(\Gamma)$ so that the solution pair (γ, y) of the FBP approximates a given boundary $\gamma_d : \Gamma \to \mathbb{R}$ and potential $y_d : \Omega^* \to \mathbb{R}$. This amounts to solving the minimization problem

$$
\min \mathcal{J}(\gamma, y, u) := \frac{1}{2}\|\gamma - \gamma_d\|^2_{L^2(\Gamma)} + \frac{1}{2}\|y - y_d\|^2_{L^2(\Omega_\gamma)} + \frac{\lambda_u}{2}\|u\|^2_{L^2(\Gamma)}, \tag{1.2a}
$$

subject to the state equations

$$
\begin{cases}
-\Delta y = 0 & \text{in } \Omega_\gamma \\
y = v & \text{on } \partial\Omega_\gamma \\
-\kappa\mathcal{H}[\gamma] + \partial_\nu y = u & \text{on } \Gamma_\gamma \\
\gamma(0) = \gamma(1) = 0,
\end{cases} \tag{1.2b}
$$

the state constraints

$$
\left|\mathrm{d}_{x_1}\gamma(x_1)\right| \le 1 \quad \text{a.e. } x_1 \in \Gamma, \tag{1.2c}
$$

with d_{x_1} being the total derivative with respect to x_1, and the control constraint

$$
u \in \mathcal{U}_{ad} \tag{1.2d}
$$

dictated by $\mathcal{U}_{ad}$, a closed ball in $L^2(\Gamma)$. Here $\lambda_u > 0$ is the stabilization parameter; v is a given data which in principle could act as a Dirichlet boundary control;

$$
\mathcal{H}[\gamma] := \mathrm{d}_{x_1}\left(\frac{\mathrm{d}_{x_1}\gamma}{\sqrt{1 + |\mathrm{d}_{x_1}\gamma|^2}}\right)
$$

is the *curvature* of γ; and $\kappa > 0$ plays the role of surface tension coefficient.

Optimal control of partial differential equations (PDEs) allows us to achieve a specific goal (1.2a) with PDE (1.2b) and other constraints (1.2c)-(1.2d) being satisfied and can be highly beneficial in practice (see [19] for more details). For example using the reverse electrowetting i.e., by applying a control to change the shape of fluid droplets, one can generate enough power to charge a cellphone [13], by mere stroll in the park. There has been various attempts to solve optimal control problems with a FBP constraint. We refer to [10, 11] for control of a two phase Stefan problem in graph formulation and [4] for the same problem in level set formulation. Paper [14] discusses optimal control of a FBP with Stokes flow. Even though problem (1.2a)-(1.2d) is relatively simple, it captures the essential features associated with surface tension effects found in more complex systems, and allows us to develop a complete second order analysis, based on [19], which is absent in the existing literature on FBP.

There are several methodologies to formulate FBP depending on the role of the free boundary. We deal with a sharp interface method, written in graph form (see Figure 1), for which the interface Γ_γ is governed by the explicit elliptic PDE

$$-\kappa \mathcal{H}\left[\gamma\right] + \partial_\nu y = u.$$

A similar approach was used in [10, 11] for the optimal control of a Stefan problem, but without the full accompanying theory developed here. Alternative approaches to treat FBP are the level set method and the diffuse interface method [6, 4].

We use a fixed domain approach to solve the optimal control free boundary problem (OC-FBP). In fact, we transform Ω_γ to $\Omega = (0,1)^2$ and Γ_γ to $\Gamma = (0,1)\times\{1\}$ (see Figure 1), at the expense of having a governing PDE with rough coefficients. This avoids dealing with shape sensitivity analysis [17, 7]. We refer to [20] for a comparison between these approaches applied to a FBP. Using operator interpolation [18] we demonstrate how to improve the existing regularity of state variables derived earlier in [16], which turns out to be instrumental to derive the second order sufficient condition. One of the challenges of an OC-FBP is dealing with possible topological changes of the domain by introducing state constraints. Our analysis provides control constraints which always enforce the state constraints i.e., we can simply treat OC-FBP as a control constrained problem.

We have organized this paper as follows. A detailed problem description on a fixed domain is given in section 2. We introduce the Lagrangian functional to formally derive the first order necessary optimality conditions in section 3. We present a rigorous justification of the Lagrangian results in the remaining sections. To this end, we introduce a control to state operator in section 4 and show that for a particular set of admissible controls it is twice Fréchet differentiable. Finally, we write the optimal control problem in its reduced form and show the existence of a control under slightly higher regularity together with second order sufficient conditions in section 5.

2. OC-FBP on Reference Domain.

For simplicity we consider the FBP (1.2b) with linearized curvature. To analyze the minimization problem (1.2), we map the physical domain Ω_γ onto the fixed reference domain $\Omega = (0,1)^2$. This results in an optimal control problem subject to PDE constraints with nonlinear coefficients depending on γ but without an explicit interface. The idea is to map the unknown domain Ω_γ onto the fixed domain $\Omega = (0,1)^2$ using the inverse of the Lipschitz map, $\Psi : \Omega \to \Omega_\gamma$, defined as

$$\Psi(x_1, x_2) = \Big(x_1, \big(1+\gamma\left(x_1\right)\big)\, x_2\Big) \in \Omega_\gamma, \quad \text{for } (x_1,x_2)\in\Omega. \tag{2.1}$$

Since γ is Lipschitz continuous according to the state constraint (1.2c), we deduce that $|\gamma| \le 1/2$ which in turn implies that Ψ is invertible. Furthermore, the inverse of Ψ is also Lipschitz. Moreover, it becomes routine to check that the Laplace equation $\Delta y = 0$ in Ω_γ and $\partial_\nu y$ on Γ_γ can be written as

$$\operatorname{div}\big(A\left[\gamma\right]\nabla y\big) = 0 \quad \text{in } \Omega, \qquad A\left[\gamma\right]\nabla y\cdot\nu\left(1+\left|\mathrm{d}_{x_1}\gamma\right|^2\right)^{-1/2} \quad \text{on } \Gamma,$$

where $\nu = [0,1]^T$, and $A : \mathring{W}^1_\infty(\Gamma) \to L^\infty(\Omega)$ is the Nemytskii operator [19, Chapter 4] defined by

$$A[\gamma] = \begin{bmatrix} 1+\gamma(x_1) & -\,\mathrm{d}_{x_1}\gamma(x_1)x_2 \\ -\,\mathrm{d}_{x_1}\gamma(x_1)x_2 & \frac{1+\left(\mathrm{d}_{x_1}\gamma(x_1)x_2\right)^2}{1+\gamma(x_1)} \end{bmatrix}. \tag{2.2}$$

It is convenient to write $A_{2,2}[\gamma]$ as $\Phi[\gamma] := \varphi\left(\gamma(x_1), \mathrm{d}_{x_1}\gamma(x_1)x_2\right)$, where $\varphi(a,b) := \left(1+b^2\right)/\left(1+a\right)$.

To simplify the exposition we make the following assumptions:

- (**A₁**) Linearized curvature: $\mathcal{H}_{lin}[\gamma] = \mathrm{d}^2_{x_1}\gamma\left(1+|\mathrm{d}_{x_1}\gamma|^2\right)^{-1/2}$.
- (**A₂**) Scaled control: $u = u\left(1+|\mathrm{d}_{x_1}\gamma|^2\right)^{-1/2}$.
- (**A₃**) Dirichlet boundary condition: we identify v on the boundary with the trace of a function $v \in W^1_p(\Omega)$.

These assumptions are not crucial. We refer to [16, Section 4] for the FBP involving $\mathcal{H}$ instead of $\mathcal{H}_{lin}$. The scaling of the control avoids unnecessarily complicating the right hand side of(2.3b) below, which would contain $u\left(1+|\mathrm{d}_{x_1}\gamma|^2\right)^{1/2}$ instead of simply u. Finally, the lifting of the Dirichlet v data is standard practice.

Under these assumptions and the application of the map Ψ, the optimal control problem (1.2) becomes

$$\min \mathcal{J}(\gamma, y, u) := \frac{1}{2}\|\gamma - \gamma_d\|^2_{L^2(\Gamma)} + \frac{\lambda_y}{2}\|y + v - y_d\|^2_{L^2(\Omega)} + \frac{\lambda_u}{2}\|u\|^2_{L^2(\Gamma)} \tag{2.3a}$$

subject to the state equations $(\gamma, y) \in \mathring{W}^1_\infty(\Gamma) \times \mathring{W}^1_p(\Omega)$

$$\begin{cases} -\operatorname{div}\left(A[\gamma]\nabla(y+v)\right) = 0 & \text{in } \Omega \\ -\kappa\,\mathrm{d}^2_{x_1}\gamma + A[\gamma]\nabla(y+v)\cdot\nu = u & \text{in } \Gamma \end{cases} \tag{2.3b}$$

together with the state constraint (1.2c) and the control constraint (1.2d).

We point out that the part of the cost functional $\mathcal{J}$ in (2.3a) corresponding to state y is defined over the fixed domain Ω instead of Ω_γ without taking into account the determinant of the Jacobian $|\mathrm{D}\Psi|$ of Ψ, because

$$\int_{\Omega_\gamma} |y + v - y_d|^2\,\mathrm{d}x_\gamma = \int_\Omega |y+v-y_d|^2 |\mathrm{D}\Psi|\,\mathrm{d}x = \int_\Omega |y+v-y_d|^2 |1+\gamma|\,\mathrm{d}x,$$

whence

$$\frac{1}{2}\int_\Omega |u + v - y_d|^2\,\mathrm{d}x \le \int_{\Omega_\gamma} |y+v-y_d|^2\,\mathrm{d}x_\gamma \le \frac{3}{2}\int_\Omega |y+v-y_d|^2\,\mathrm{d}x,$$

where we have used $|\gamma| \le \frac{1}{2}$. Therefore, we define $\mathcal{J}$ over Ω without $|\mathrm{D}\Psi|$.

In order to derive the first and second order optimality conditions in later sections, we need to compute the first and second order directional derivatives of A, which in turn requires computing the directional derivative of the Nemytskii operator Φ defined above. To simplify notation, we drop the evaluation of γ and $\mathrm{d}_{x_1}\gamma$ at x_1. The derivative of Φ in the direction h at $(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)$ is given by

$$\mathrm{D}\Phi[\gamma]\langle h\rangle = \partial_a\varphi\left(\gamma, x_2\,\mathrm{d}_{x_1}\gamma\right)h + \partial_b\varphi\left(\gamma, x_2\,\mathrm{d}_{x_1}\gamma\right)x_2\,\mathrm{d}_{x_1}h$$

where

$$\nabla\varphi(a,b) := \begin{bmatrix}\partial_a\varphi(a,b) & \partial_b\varphi(a,b)\end{bmatrix} = \begin{bmatrix}-\frac{1+b^2}{(1+a)^2} & \frac{2b}{1+a}\end{bmatrix}.$$

Furthermore, we obtain the following representation for $\mathrm{D}A$ in terms of h and h'

$$\begin{aligned}\mathrm{D}A\,[\gamma]\,\langle h\rangle &:= A_1\,[\gamma]\,h + A_2\,[\gamma]\,\mathrm{d}_{x_1}h \\ &= \begin{bmatrix}1 & 0 \\ 0 & \partial_a\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\end{bmatrix} h + \begin{bmatrix}0 & -x_2 \\ -x_2 & x_2\partial_b\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\end{bmatrix}\mathrm{d}_{x_1}h, \qquad (2.4)\end{aligned}$$

whence the remainder $\mathfrak{R}_A\,[\gamma, h]$ at γ in the direction h reads

$$\mathfrak{R}_A\,[\gamma, h] := A\,[\gamma + h] - A\,[\gamma] - \mathrm{D}A\,[\gamma]\,\langle h\rangle \qquad (2.5a)$$

and

$$\lim_{|h|_{W^1_\infty(\Gamma)}\to 0} \frac{\left\|\mathfrak{R}_A\,[\gamma, h]\right\|_{L^\infty(\Gamma)}}{|h|_{W^1_\infty(\Gamma)}} = 0. \qquad (2.5b)$$

The Hessian of φ is

$$\nabla^2\varphi(a,b) = \begin{bmatrix}\partial_a^2\varphi(a,b) & \partial_{ab}\varphi(a,b) \\ \partial_{ba}\varphi(a,b) & \partial_b^2\varphi(a,b)\end{bmatrix} = 2\begin{bmatrix}\frac{1+b^2}{(1+a)^3} & \frac{-b}{(1+a)^2} \\ \frac{-b}{(1+a)^2} & \frac{1}{1+a}\end{bmatrix}.$$

The second order derivative of Φ in the direction h_1 followed by h_2 evaluated at $(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)$ is

$$\begin{aligned}\mathrm{D}^2\,\Phi\,[\gamma]\,\langle h_2, h_1\rangle &= \partial_a^2\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\,h_2h_1 + \partial_{ab}\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\,x_2h_2\,\mathrm{d}_{x_1}h_1 \\ &\quad + \partial_{ab}\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\,x_2\,\mathrm{d}_{x_1}h_2h_1 + \partial_b^2\varphi\,(\gamma, x_2\,\mathrm{d}_{x_1}\gamma)\,x_2^2\,\mathrm{d}_{x_1}h_2\,\mathrm{d}_{x_1}h_1.\end{aligned}$$

Finally, we obtain the following representation for D^2A in terms of h_1 and h_2

$$\mathrm{D}^2A\,[\gamma]\,\langle h_2, h_1\rangle = \begin{bmatrix}0 & 0 \\ 0 & \mathrm{D}^2\,\Phi\,[\gamma]\,\langle h_2, h_1\rangle\end{bmatrix}, \qquad (2.6)$$

whence the remainder $\mathfrak{R}_{\mathrm{D}A}\,[\gamma, h_1, h_2]$ at γ reads

$$\mathfrak{R}_{\mathrm{D}A}\,[\gamma, h_1, h_2] := \mathrm{D}A\,[\gamma + h_2]\,\langle h_1\rangle - \mathrm{D}A\,[\gamma]\,\langle h_1\rangle - \mathrm{D}^2A\,[\gamma]\,\langle h_1, h_2\rangle, \qquad (2.7a)$$

and

$$\lim_{\substack{|h_1|_{W^1_\infty(\Gamma)}\to 0 \\ |h_2|_{W^1_\infty(\Gamma)}\to 0}} \frac{\left\|\mathfrak{R}_{\mathrm{D}A}\,[\gamma, h_1, h_2]\right\|_{L^\infty(\Omega)}}{|h_1|_{W^1_\infty(\Gamma)}|h_2|_{W^1_\infty(\Gamma)}} = 0. \qquad (2.7b)$$

Proposition 2.1 (bounds on A). *Applying the state constraint* (1.2c) *to* (2.2), (2.4) *and* (2.6) *implies the existence of a positive constant $C_A < \infty$ such that*

$$\begin{aligned}\left\|A\,[\gamma]\right\|_{L^\infty(\Omega)} &+ \sup_{|h|_{W^1_\infty(\Omega)}=1}\left\|\mathrm{D}A\,[\gamma]\,\langle h\rangle\right\|_{L^\infty(\Omega)} \\ &+ \sup_{\substack{|h_1|_{W^1_\infty(\Omega)}=1 \\ |h_2|_{W^1_\infty(\Omega)}=1}}\left\|\mathrm{D}^2A\,[\gamma]\,\langle h_1, h_2\rangle\right\|_{L^\infty(\Omega)} \le C_A. \qquad (2.8)\end{aligned}$$

3. Lagrangian Formulation. In this section we formally derive, ignoring the state constraint (1.2c), the first order necessary optimality conditions using the Lagrangian approach described in [19]. For a rigorous analysis of the existence of Lagrange multipliers in Banach spaces we refer to [23]. It is well known that for a convex optimal control problem with linear constraints, the first order necessary optimality conditions are also sufficient conditions [19, Lemma 2.21]. However, despite linearizing the curvature via assumption ($\mathbf{A_1}$), the state equations (2.3b) are still highly nonlinear, and the optimization nonconvex. We will derive the second order sufficient optimality conditions in section 5.

Let s, r denote the adjoint variables corresponding to states γ, y respectively, and p, q be conjugate indices i.e., $1/p + 1/q = 1$ with $p > 2$. Then the Lagrangian functional is given by

$$\begin{aligned}\mathcal{L}(\gamma, y, u, r, s) := \mathcal{J}(\gamma, y, u) &+ \int_\Omega \operatorname{div}\big(A[\gamma]\nabla(y+v)\big)\, r \,\mathrm{d}x \\ &+ \int_\Gamma \Big(\kappa\, \mathrm{d}_{x_1}^2 \gamma - A[\gamma]\nabla(y+v)\cdot\nu + u\Big)\, s\,\mathrm{d}\sigma.\end{aligned} \tag{3.1}$$

Additionally, if $(\bar\gamma, \bar y, \bar u, \bar r, \bar s)$ is a critical point for the Lagrangian $\mathcal{L}$, then the first order necessary optimality conditions are

$$\Big\langle \mathrm{D}_{\{\gamma,y,r,s\}}\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), h\Big\rangle = 0 \quad \forall h \in \Big\{\mathring{W}^1_\infty(\Gamma), \mathring{W}^1_p(\Omega), \mathring{W}^1_q(\Omega), \mathring{W}^1_1(\Gamma)\Big\}, \tag{3.2a}$$

$$\big\langle \mathrm{D}_u\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), u - \bar u\big\rangle \geq 0 \quad \forall u \in \mathcal{U}_{ad}, \tag{3.2b}$$

where $\{\}$ denotes a list, e.g. $\big\langle \mathrm{D}_s\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), h\big\rangle = 0$ for all $h \in \mathring{W}^1_1(\Gamma)$. Therefore, computing $(\bar\gamma, \bar y, \bar u, \bar r, \bar s)$ requires solving the nonlinear system (3.2). In practice this can be realized using techniques described in [3, 12, 19]. To solve variational inequalities of first and second kind we refer to [9] for relaxation and augmented Lagrangian techniques and to [5] for semi-smooth Newton methods, and references therein. The remainder of this section is devoted to the derivation of the equations satisfied by $(\bar\gamma, \bar y, \bar u, \bar r, \bar s)$ using the nonlinear system above.

Since $\Big\langle \mathrm{D}_{\{s,r\}}\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), h\Big\rangle = 0$ implies that $(\bar\gamma, \bar y)$ solves the state equations (2.3b), we focus on the adjoint equations $\Big\langle \mathrm{D}_{\{\gamma,y\}}\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), h\Big\rangle = 0$. Using Green's theorem and assuming smoothness, the Lagrangian $\mathcal{L}$ can be rewritten as:

$$\begin{aligned}\mathcal{L}(y, \gamma, u, r, s) = \mathcal{J}(\gamma, y, u) &+ \int_\Omega \operatorname{div}\big(A[\gamma]\nabla r\big)(y+v)\,\mathrm{d}x \\ &+ \int_{\partial\Omega} \big(rA[\gamma]\nabla(y+v) - (y+v)A[\gamma]\nabla r\big)\cdot\nu\,\mathrm{d}\sigma \\ &+ \int_\Gamma \Big(\kappa\gamma\, \mathrm{d}_{x_1}^2 s - sA[\gamma]\nabla(y+v)\cdot\nu + us\Big)\,\mathrm{d}\sigma + \kappa\left(\mathrm{d}_{x_1}\gamma s - \gamma\,\mathrm{d}_{x_1} s\right)\Big|_0^1.\end{aligned} \tag{3.3}$$

Imposing $\big\langle \mathrm{D}_y\, \mathcal{L}(\bar\gamma, \bar y, \bar u, \bar r, \bar s), h\big\rangle = 0$ to (3.3) implies that for every $h \in \mathring{W}^1_p(\Omega)$

$$\begin{aligned}-\int_\Omega \operatorname{div}\big(A[\bar\gamma]\nabla\bar r\big)\, h\,\mathrm{d}x = \lambda_y \int_\Omega (\bar y + v - y_d)\, h\,\mathrm{d}x \\ + \int_{\partial\Omega} \bar r A[\bar\gamma]\nabla h\cdot\nu\,\mathrm{d}\sigma - \int_\Gamma \bar s A[\bar\gamma]\nabla h\cdot\nu\,\mathrm{d}\sigma.\end{aligned} \tag{3.4a}$$

Next, without loss of generality ($C_0^\infty(\Omega)$ is dense in $\mathring{W}_p^1(\Omega)$), we obtain

$$-\int_\Omega \operatorname{div}\left(A[\bar{\gamma}]\nabla\bar{r}\right) h\,\mathrm{d}x = \lambda_y \int_\Omega (\bar{y}+v-y_d)\,h\,\mathrm{d}x \qquad \forall h\in C_0^\infty(\Omega), \tag{3.4b}$$

whereas, using that $A[\gamma]\nabla h\cdot\nu$ can be chosen arbitrarily on $\partial\Omega$ we deduce from (3.4a) and (3.4b) that

$$\bar{r}-\bar{s}|_\Gamma = 0, \qquad \bar{r}|_\Sigma = 0. \tag{3.4c}$$

In view of (3.4b-c), the strong form of the boundary value problem for $\bar{r}$ is: seek $\bar{r}\in W_q^1(\Omega)$ such that

$$\begin{cases} -\operatorname{div}\left(A[\bar{\gamma}]\,\bar{r}\right) = \lambda_y(\bar{y}+v-y_d) & \text{in } \Omega \\ \bar{r}=\bar{s} & \text{on } \Gamma \\ \bar{r}=0 & \text{on } \Sigma. \end{cases} \tag{3.5}$$

Next we employ the same technique to obtain the equations for the second adjoint variable $\bar{s}$: we impose $\left\langle \mathrm{D}_\gamma\,\mathcal{L}(\bar{\gamma},\bar{y},\bar{u},\bar{r},\bar{s}),h\right\rangle = 0$ to (3.3) and make use of the boundary conditions in (3.5) to obtain for every $h\in\mathring{W}_\infty^1(\Gamma)$

$$-\int_\Gamma \kappa\,\mathrm{d}_{x_1}^2 sh\,\mathrm{d}\sigma = \int_\Gamma (\bar{\gamma}-\gamma_d)\,h\,\mathrm{d}\sigma - \int_\Omega \mathrm{D}A[\bar{\gamma}]\langle h\rangle\nabla(\bar{y}+v)\cdot\nabla\bar{r}\,\mathrm{d}x.$$

Therefore, the strong form of the boundary value problem for $\bar{s}$ is: seek $\bar{s}\in\mathring{W}_1^1(\Gamma)$

$$\begin{cases} -\kappa\,\mathrm{d}_{x_1}^2\bar{s} = (\bar{\gamma}-\gamma_d) - \displaystyle\int_0^1 A_1[\bar{\gamma}]\nabla(\bar{y}+v)\cdot\nabla\bar{r}\,\mathrm{d}x_2 \\ \qquad\qquad + \mathrm{d}_{x_1}\left(\displaystyle\int_0^1 A_2[\bar{\gamma}]\nabla(\bar{y}+v)\cdot\nabla\bar{r}\,\mathrm{d}x_2\right) \quad \text{in } \Gamma \\ \bar{s}(0)=\bar{s}(1)=0, \end{cases} \tag{3.6}$$

where A_1, A_2 denote the representation of DA given in (2.4). We note that the integrals on the right hand side of (3.6) correspond to integration in x_2 (vertical) direction.

Finally, (3.2b) implies

$$\langle\lambda_u\bar{u}+\bar{s},u-\bar{u}\rangle_{L^2(\Gamma)\times L^2(\Gamma)} \geq 0, \qquad \forall u\in\mathcal{U}_{ad}. \tag{3.7}$$

To summarize, the solution $(\bar{\gamma},\bar{y},\bar{u},\bar{r},\bar{s})$ to the first order optimality system (3.2) satisfies (2.3b), (3.5), (3.6) and (3.7). We stress that the formal approach presented in this section is very systematic and highly useful even though it is not clear at the moment how to show the existence and (local) uniqueness of the optimal control $\bar{u}$. A rigorous analysis will be developed in the next two sections.

4. The Control to State Map G_v.

Let G_v denote the nonlinear map

$$\begin{array}{cccc} G_v & : & \mathcal{U}_{ad} & \longrightarrow & \mathbb{W}^1 \\ & & u & \longmapsto & (\gamma,y) \end{array}, \tag{4.1}$$

where $\mathbb{W}^1 := \mathring{W}^1_\infty(\Gamma) \times \mathring{W}^1_p(\Omega)$, (γ, y) solves (2.3b), and the subscript on G_v denotes dependence on a fixed and non-trivial $v \in W^1_p(\Omega)$. Our goal is to show the existence of a control, derive the first order necessary and second order sufficient optimality conditions within the realm of a rigorous mathematical framework. The first order optimality conditions requires to show that G_v is Fréchet differentiable (subsection 4.3) and the second order conditions require G_v to be twice Fréchet differentiable (subsection 4.4).

The steps described above are standard for PDE-constrained optimization in fixed domains[19], but our analysis for the linearized curvature OC-FBP is novel. The novelty resides in the highly nonlinear structure of the underlying FBP, which is posed in a pair of Banach spaces one being non-reflexive, and yet we deal with *minimal regularity*. A number of other control problems for FBP fall under a similar functional framework [1, 2], but their theory is not as complete and conclusive as ours. This appears to be an area of intense current research.

The first step in this voyage is to show that there exists a unique weak solution to (2.3b), which implies that G_v is a well defined one-to-one nonlinear operator. In fact, it is known [16] that for $u = 0$ and v small, a fixed point argument asserts the existence and uniqueness of a weak solution (γ, y) in $\mathbb{W}^1$ to (2.3b). We further extend this analysis to the case where $u \neq 0$. This gives us an admissible ball $\mathcal{U}_{ad} \subset L^2(\Gamma)$ where we can show the existence of solution to (2.3b). Furthermore, this set guarantees that the state constraints in (1.2c) are always satisfied, thereby leading to a standard control constrained problem without state constraints.

4.1. Well-posedness of the State System (2.3b)**.** The weak form of the system (2.3b) is: find $(\gamma, y) \in \mathbb{W}^1$ such that

$$\begin{cases} \mathcal{B}_\Omega\left[y + v, z; A[\gamma]\right] = 0 & \forall z \in \mathring{W}^1_q(\Omega) \\ \mathcal{B}_\Gamma[\gamma, \zeta] + \mathcal{B}_\Omega\left[y + v, E\zeta; A[\gamma]\right] = \langle u, \zeta\rangle_{W^{-1}_\infty(\Gamma)\times\mathring{W}^1_1(\Gamma)} & \forall \zeta \in \mathring{W}^1_1(\Gamma), \end{cases} \tag{4.2}$$

where $\mathcal{B}_\Gamma : \mathring{W}^1_\infty(\Gamma) \times \mathring{W}^1_1(\Gamma) \to \mathbb{R}$, $\mathcal{B}_\Omega : \mathring{W}^1_p(\Omega) \times \mathring{W}^1_q(\Omega) \to \mathbb{R}$ are defined by

$$\begin{aligned} \mathcal{B}_\Gamma[\gamma, \zeta] &:= \kappa \int_\Gamma \mathrm{d}_{x_1}\gamma(x_1)\mathrm{d}_{x_1}\zeta(x_1)\,\mathrm{d}x_1, \\ \mathcal{B}_\Omega\left[y, z; A[\gamma]\right] &:= \int_\Omega A[\gamma]\nabla y \cdot \nabla z\,\mathrm{d}x. \end{aligned} \tag{4.3}$$

Furthermore, $E : \mathring{W}^1_1(\Gamma) \to W^1_q(\Omega)$, $q < 2$ denotes a continuous extension such that $E\zeta|_\Gamma = \zeta$, $E\zeta|_\Sigma = 0$ (cf. [16, Lemma 2]). In particular, this implies that there exists a constant $C_E \geq 1$ such that

$$|E\zeta|_{W^1_q(\Omega)} \leq C_E|\zeta|_{W^1_1(\Gamma)}, \quad \forall \zeta \in \mathring{W}^1_1(\Gamma). \tag{4.4}$$

Moreover, for $u \in \mathcal{U}_{ad} \subset L^2(\Gamma)$ we can write

$$\langle u, \zeta\rangle_{W^{-1}_\infty(\Gamma)\times\mathring{W}^1_1(\Gamma)} = -\left\langle \int_0^{x_1} u, \mathrm{d}_{x_1}\zeta \right\rangle_{L^\infty(\Gamma)\times L^1(\Gamma)}. \tag{4.5}$$

In view of the regularity of u and ζ, the duality pairing in (4.5) reduces to $\int_0^1 u\zeta$. This also enables us to deduce that for $u \in L^2(\Gamma)$

$$\|u\|_{W^{-1}_\infty(\Gamma)} \leq \left\| \int_0^{x_1} u \right\|_{L^\infty(\Gamma)} \leq \|u\|_{L^2(\Gamma)}. \tag{4.6}$$

We will make use of these two facts repeatedly throughout the rest of the paper.

Proposition 4.1 (inf-sup conditions). *The following conditions hold for the bilinear forms* $\mathcal{B}_\Gamma\left[\cdot,\cdot\right]$ *and* $\mathcal{B}_\Omega\left[\cdot,\cdot;A\left[\gamma\right]\right]$ *defined in* (4.3) *:*

(i) $\mathcal{B}_\Gamma\left[\cdot,\cdot\right]$ *is continuous and there exists a constant* $\alpha > 0$ *such that for every* $\gamma \in \mathring{W}^1_\infty\left(\Gamma\right)$ *and* $s \in \mathring{W}^1_1\left(\Gamma\right)$

$$|\gamma|_{W^1_\infty(\Gamma)} \leq \alpha \sup_{0 \neq \zeta \in \mathring{W}^1_1(\Gamma)} \frac{\mathcal{B}_\Gamma\left[\gamma,\zeta\right]}{|\zeta|_{W^1_1(\Gamma)}}, \tag{4.7a}$$

$$|s|_{W^1_1(\Gamma)} \leq \alpha \sup_{0 \neq \zeta \in \mathring{W}^1_\infty(\Gamma)} \frac{\mathcal{B}_\Gamma\left[\zeta,s\right]}{|\zeta|_{W^1_\infty(\Gamma)}}. \tag{4.7b}$$

(ii) If $\gamma \in \mathring{W}^1_\infty\left(\Gamma\right)$ *satisfies* (1.2c)*, then* $\mathcal{B}_\Omega\left[\cdot,\cdot;A\left[\gamma\right]\right]$ *is continuous and there exist constants* P, Q *with* $Q < 2 < P$ *and* $\beta > 0$*, such that for* $p \in (Q, P)$

$$|y|_{W^1_p(\Omega)} \leq \beta \sup_{0 \neq z \in \mathring{W}^1_q(\Omega)} \frac{\mathcal{B}_\Omega\left[y,z;A\left[\gamma\right]\right]}{|z|_{W^1_q(\Omega)}}. \tag{4.8}$$

Proof. For (4.7a) and (4.8) we refer to [16, Proposition 2.2-2.3] for a proof. For (4.7b) we proceed as follows: applying the definition of the L^1-norm and the homogeneous Dirichlet values of s, we obtain

$$\begin{aligned} |s|_{W^1_1(\Gamma)} &= \int_0^1 |\mathrm{d}_{x_1} s| = \int_0^1 \mathrm{sgn}(\mathrm{d}_{x_1} s)\, \mathrm{d}_{x_1} s \\ &= \int_0^1 \underbrace{\left(\mathrm{sgn}(\mathrm{d}_{x_1} s) - \int_0^1 \mathrm{sgn}(\mathrm{d}_{x_1} s)\right)}_{=\mathrm{d}_{x_1}\zeta} \mathrm{d}_{x_1} s = \frac{1}{\kappa}\mathcal{B}_\Gamma\left[\zeta,s\right], \end{aligned}$$

where $\zeta(x_1) = \int_0^{x_1}\left(\mathrm{sgn}(\mathrm{d}_{x_1} s) - \int_0^1 \mathrm{sgn}(\mathrm{d}_{x_1} s)\right) \in \mathring{W}^1_\infty\left(\Gamma\right)$. Estimate (4.7b) follows by noting that $|\zeta|_{W^1_\infty(\Gamma)} \leq 2$, and taking the sup over every $\zeta \in \mathring{W}^1_\infty\left(\Gamma\right)$. ◻

The following lemma demonstrates how one can improve the integrability index of a solution to a PDE obtained by standard methods.

Lemma 4.2 (improved integrability). *Let* Ω *be an open Lipschitz bounded domain of* $\mathbb{R}^d$ *and* $\mathcal{B} : \mathring{W}^1_\infty\left(\Omega\right) \times \mathring{W}^1_1\left(\Omega\right) \to \mathbb{R}$ *be a continuous bilinear form. Furthermore, suppose that*

(i) there exists $\alpha > 0$ *such that*

$$|\chi|_{W^1_\infty(\Omega)} \leq \alpha \sup_{0 \neq \psi \in \mathring{W}^1_1(\Omega)} \frac{\mathcal{B}\left[\chi,\psi\right]}{|\psi|_{W^1_1(\Omega)}} \qquad \forall \chi \in \mathring{W}^1_\infty\left(\Omega\right), \tag{4.9}$$

(ii) and $\mathcal{B}$ *is continuous and coercive in* $\mathring{W}^1_2\left(\Omega\right)$*.*

Then for every $F \in \mathring{W}^1_1\left(\Omega\right)^*$*, there exists a unique* $\chi \in \mathring{W}^1_\infty\left(\Omega\right)$ *such that*

$$\mathcal{B}\left[\chi,\psi\right] = F(\psi) \text{ for all } \psi \in \mathring{W}^1_1\left(\Omega\right) \text{ and } |\chi|_{W^1_\infty(\Omega)} \leq \alpha \|F\|_{W^1_1(\Omega)^*}. \tag{4.10}$$

Proof. Since $\mathring{W}^1_2\left(\Omega\right) \subset \mathring{W}^1_1\left(\Omega\right)$, it follows that $F \in \mathring{W}^1_1\left(\Omega\right)^* \subset \mathring{W}^1_2\left(\Omega\right)^*$. Lax-Milgram guarantees the existence and uniqueness of $\chi \in \mathring{W}^1_2\left(\Omega\right)$ such that $\mathcal{B}\left[\chi,\psi\right] = F(\psi)$ for all $\psi \in \mathring{W}^1_2\left(\Omega\right)$.

Next, we extend $\mathcal{B}[\chi,\cdot]$ as a linear functional on $\mathring{W}^1_1(\Omega)$. To this end, let $\{\psi_n\} \subset \mathring{W}^1_2(\Omega)$ be a Cauchy sequence in the $\mathring{W}^1_1(\Omega)$-norm. It immediately follows that $\big\{\mathcal{B}[\chi,\psi_n]\big\}$ is also Cauchy in $\mathbb{R}$, i.e.

$$\big|\mathcal{B}[\chi,\psi_n-\psi_m]\big| = \big|F(\psi_n-\psi_m)\big| \leq \|F\|_{\mathring{W}^1_1(\Omega)^*} |\psi_n-\psi_m|_{W^1_1(\Omega)}.$$

Finally, through density of $\mathring{W}^1_2(\Omega)$ in $\mathring{W}^1_1(\Omega)$, not only do we obtain $\psi_n \to \psi \in \mathring{W}^1_1(\Omega)$, but also

$$\mathcal{B}[\chi,\psi] := \lim_{n\to\infty} \mathcal{B}[\chi,\psi_n] = \lim_{n\to\infty} F(\psi_n) = F(\psi).$$

The estimate for $|\chi|_{W^1_\infty(\Gamma)}$ follows by (4.9). □

4.1.1. First order regularity. Now we are ready to prove that there exists a unique solution to (4.2) with $\mathbb{W}^1$-regularity. Since the system (4.2) is nonlinear we will obtain this result by applying the Banach fixed point theorem combined with a smallness assumption on a non-trivial v. To this end, we let $2 < p < P$ and equip the space $\mathbb{W}^1 = \mathring{W}^1_\infty(\Gamma) \times \mathring{W}^1_p(\Omega)$ with the equivalent norm

$$\big\|(\gamma,y)\big\|_{\mathbb{W}^1} := (1+\beta C_A)|v|_{W^1_p(\Omega)}|\gamma|_{W^1_\infty(\Gamma)} + |y|_{W^1_p(\Omega)}, \tag{4.11}$$

where C_A and β are given in (2.8) and (4.8), and define the closed (convex) ball

$$\mathbb{B}_v := \Big\{(\gamma,y) \in \mathbb{W}^1 : |y|_{W^1_p(\Omega)} \leq \beta C_A |v|_{W^1_p(\Omega)},\, |\gamma|_{W^1_\infty(\Gamma)} \leq 1\Big\}. \tag{4.12}$$

Furthermore, consider the operator $T : \mathbb{B}_v \to \mathbb{W}^1$ defined as

$$T(\gamma,y) := \big(T_1(\gamma,y), T_2(\gamma,y)\big) = (\widetilde{\gamma},\widetilde{y}) \qquad \forall\,(\gamma,y) \in \mathbb{B}_v, \tag{4.13}$$

where $\widetilde{\gamma} = T_1(\gamma,y) \in \mathring{W}^1_\infty(\Gamma)$ satisfies for every $\zeta \in \mathring{W}^1_1(\Gamma)$

$$\mathcal{B}_\Gamma[\widetilde{\gamma},\zeta] = -\mathcal{B}_\Omega\big[y+v, E\zeta; A[\gamma]\big] + \langle u,\zeta\rangle_{W^{-1}_\infty(\Gamma)\times\mathring{W}^1_1(\Gamma)}, \tag{4.14}$$

and $\widetilde{y} = T_2(\gamma,y) \in \mathring{W}^1_p(\Omega)$ satisfies for every $z \in \mathring{W}^1_q(\Omega)$

$$\mathcal{B}_\Omega\Big[\widetilde{y}+v, z; A\big[T_1(\gamma,y)\big]\Big] = 0. \tag{4.15}$$

With these definitions at hand we proceed to find conditions under which T not only maps $\mathbb{B}_v$ into itself but is in fact a contraction in $\mathbb{B}_v$.

Lemma 4.3 (range of T). *Let T_1 and T_2 be the operators defined in* (4.14) *and* (4.15)*, and C_A and C_E be the constants defined in* (2.8) *and* (4.4)*. Furthermore, suppose there exists $\theta_1 \in \big(\beta C_A/(1+\beta C_A), 1\big)$ such that*

$$|v|_{W^1_p(\Omega)} \leq (1-\theta_1)\big(\alpha C_E C_A(1+\beta C_A)\big)^{-1}. \tag{4.16}$$

If $u \in L^2(\Gamma)$ with $\|u\|_{L^2(\Gamma)} \leq \theta_1/\alpha$, then the range of T is contained in $\mathbb{B}_v$.

Proof. Let $(\gamma,y) \in \mathbb{B}_v$ be fixed but arbitrary. First we rely on Lemma 4.2 to show the well-posedness of T_1. Since it is straight-forward to check that $\mathcal{B}_\Gamma$ is continuous and coercive in $\mathring{W}^1_2(\Gamma)$, we only need to show the regularity of the forcing term in

(4.14). If we define $F(\zeta) := -\mathcal{B}_\Omega\left[y+v, E\zeta; A\left[\gamma\right]\right] + \langle u, \zeta\rangle$ and use (2.8), (4.4) and (4.12) we find that

$$\begin{aligned} \big|F(\zeta)\big| &\le C_A\left(|y|_{W_p^1(\Omega)} + |v|_{W_p^1(\Omega)}\right)|E\zeta|_{W_q^1(\Omega)} + \|u\|_{L^2(\Gamma)}|\zeta|_{W_1^1(\Gamma)} \\ &\le \left(C_E C_A\left(1+\beta C_A\right)|v|_{W_p^1(\Omega)} + \|u\|_{L^2(\Gamma)}\right)|\zeta|_{W_1^1(\Gamma)}\,, \end{aligned} \tag{4.17}$$

whence $F \in \mathring{W}_1^1\left(\Gamma\right)^*$ and we conclude from (4.10) that

$$|\widetilde{\gamma}|_{W_\infty^1(\Gamma)} \le \alpha\left(C_E C_A\left(1+\beta C_A\right)|v|_{W_p^1(\Omega)} + \|u\|_{L^2(\Gamma)}\right) \le (1-\theta_1) + \theta_1 = 1.$$

The well-posedness of T_2 follows by Proposition 4.1 and the Banach-Nečas theorem for reflexive Banach spaces [8, Theorem 2.6]. Applying (4.8) we obtain

$$|\widetilde{y}|_{W_p^1(\Omega)} \le \beta C_A |v|_{W_p^1(\Omega)}\,.$$

Since (γ, y) is arbitrary, we conclude that the range of T is contained in $\mathbb{B}_v$. ☐

Definition 4.4 (admissible control set). *Let θ_1 be as defined in Lemma 4.3. The admissible set of controls $\mathcal{U}_{ad}$ is the (nontrivial) closed ball*

$$\mathcal{U}_{ad} := \left\{u \in L^2\left(\Gamma\right)\,:\, \|u\|_{L^2(\Gamma)} \le \theta_1/\alpha\right\}. \tag{4.18}$$

Theorem 4.5 (T is a contraction). *Let the assumptions of Lemma 4.3 hold and suppose further that there exists a $\theta_2 \in (0,1)$ such that*

$$|v|_{W_p^1(\Omega)} \le (1-\theta_2)\left(\alpha C_E C_A\left(1+\beta C_A\right)^2\right)^{-1}. \tag{4.19}$$

Then, the map T defined in (4.13) *is a contraction in $\mathbb{B}_v$ with constant $1-\theta_2$.*

Proof. Consider $(\gamma_1, y_1), (\gamma_2, y_2) \in \mathbb{B}_v$ such that $(\gamma_1, y_1) \neq (\gamma_2, y_2)$. Using (4.13) we have that $T\left(\gamma_i, y_i\right) = \left(\widetilde{\gamma}_i, \widetilde{y}_i\right)$ solves (4.14) and (4.15) for $i = 1, 2$. Therefore, combining Proposition 4.1 (i) and Lemma 4.3 with (4.19) implies

$$\begin{aligned} |\widetilde{\gamma}_1 - \widetilde{\gamma}_2|_{W_\infty^1(\Gamma)} &\le \alpha \sup_{|\zeta|_{W_1^1(\Gamma)}=1} \mathcal{B}_\Gamma\left[\widetilde{\gamma}_1 - \widetilde{\gamma}_2, \zeta\right] \\ &= \alpha \sup_{|\zeta|_{W_1^1(\Gamma)}=1} \mathcal{B}_\Omega\left[y_2 - y_1, E\zeta; A\left[\gamma_1\right]\right] + \mathcal{B}_\Omega\left[y_2 + v, E\zeta; A\left[\gamma_2\right] - A\left[\gamma_1\right]\right] \\ &\le \alpha C_E C_A\left(|y_1 - y_2|_{W_p^1(\Omega)} + (1+\beta C_A)|v|_{W_p^1(\Omega)}|\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)}\right) \\ &= \alpha C_E C_A \big\|(\gamma_1 - \gamma_2, y_1 - y_2)\big\|_{\mathbb{W}^1}\,. \end{aligned} \tag{4.20}$$

Similarly, Proposition 4.1 (ii) in conjunction with (4.15) implies

$$\begin{aligned} |\widetilde{y}_1 - \widetilde{y}_2|_{W_p^1(\Omega)} &\le \beta \sup_{|z|_{W_q^1(\Omega)}=1} \mathcal{B}_\Omega\left[\widetilde{y}_1 - \widetilde{y}_2, z; A\left[\widetilde{\gamma}_1\right]\right] \\ &= \beta \sup_{|z|_{W_q^1(\Omega)}=1} \mathcal{B}_\Omega\left[\widetilde{y}_2 + v, z; A\left[\widetilde{\gamma}_2\right] - A\left[\widetilde{\gamma}_1\right]\right] \qquad (4.21) \\ &\le \beta C_A |\widetilde{y}_2 + v|_{W_p^1(\Omega)}|\widetilde{\gamma}_1 - \widetilde{\gamma}_2|_{W_\infty^1(\Gamma)} \\ &\le \beta C_A\left(1+\beta C_A\right)|v|_{W_p^1(\Omega)}|\widetilde{\gamma}_1 - \widetilde{\gamma}_2|_{W_\infty^1(\Gamma)}\,. \end{aligned}$$

Finally, (4.20) and (4.21) imply

$$\begin{aligned}\big\|(\widetilde{\gamma}_1,\widetilde{y}_1)-(\widetilde{\gamma}_2,\widetilde{y}_2)\big\|_{\mathbb{W}^1} &\leq (1+\beta C_A)^2|v|_{W^1_p(\Omega)}|\widetilde{\gamma}_1-\widetilde{\gamma}_2|_{W^1_\infty(\Gamma)}\\ &\leq \alpha C_E C_A\,(1+\beta C_A)^2|v|_{W^1_p(\Omega)}\big\|(\gamma_1,y_1)-(\gamma_2,y_2)\big\|_{\mathbb{W}^1}\\ &\leq (1-\theta_2)\big\|(\gamma_1,y_1)-(\gamma_2,y_2)\big\|_{\mathbb{W}^1}\,,\end{aligned}$$

where the last inequality follows from (4.19). Since $\theta_2\in(0,1)$, T is a contraction with constant $1-\theta_2$ as asserted. ☐

Corollary 4.6 (well-posedness of state system). *For every $u\in\mathcal{U}_{ad}$, the closed ball of Definition 4.4, and v satisfying* (4.16) *and* (4.19)*, there exists a unique solution $(\gamma,y)\in\mathbb{W}^1$ to the state equations* (4.2)*. This further implies that G_v in* (4.1) *is a well defined, one-to-one, nonlinear operator.*

Proof. Let $u\in\mathcal{U}_{ad}$ be fixed but arbitrary. It now follows that T is a contraction in the closed convex set $\mathbb{B}_v$ (cf. Theorem 4.5) and applying the Banach fixed point theorem we obtain a unique $(\gamma,y)\in\mathbb{B}_v$ such that $T(\gamma,y)=(\gamma,y)$. In view of (4.14) and (4.15), this is equivalent to saying that (γ,y) is the weak solution to the FBP (4.2), i.e. $G_v(u)=(\gamma,y)$. ☐

4.1.2. Enhanced Regularity of γ. Corollary 4.6 implies the existence and uniqueness of a solution (γ,y) to (2.3b) with $\mathbb{W}^1$-regularity, provided $u\in\mathcal{U}_{ad}$ and v satisfies (4.16) and (4.19). That is, we only have one weak derivative for γ and y. In the sequel we will show that the solution $(\gamma,y)=G_v(u)$ is slightly more regular *without* any extra restrictions on u or v. More specifically, we will show that

$$\gamma\in W^{1+1/q}_p(\Gamma)\cap\mathring{W}^1_\infty(\Gamma). \tag{4.22}$$

The importance of this result will be evident in subsection 5.1 where we show the existence of an optimal control. Despite its importance, the proof is rather simple.

Let $(\gamma,y)\in\mathbb{W}^1$ be a weak solution to (4.2). Function γ satisfies

$$-\kappa\,\mathrm{d}^2_{x_1}\gamma=-A[\gamma]\nabla(y+v)\cdot\nu+u=:f,$$

in the sense of distributions. If we assume, for the moment, that $f\in W^{-1/p}_p(\Gamma)$, then $\mathrm{d}^2_{x_1}\gamma\in W^{-1/p}_p(\Gamma)$. This directly implies $\gamma\in W^{-1/p+2}_p(\Gamma)$, i.e. $\gamma\in W^{1+1/q}_p(\Gamma)$ as desired. Thus, it remains to show that f is in $W^{-1/p}_p(\Gamma)$ as suggested. Since $u\in L^2(\Gamma)$, we just need to deal with the first term.

If $\phi\in\mathring{W}^{1/p}_q(\Gamma)$ then it can be seen as the trace of a function $E\phi\in W^1_q(\Omega)$ such that $E\phi|_\Gamma=\phi$, $E\phi|_\Sigma=0$, and $\|E\phi\|_{W^1_q(\Omega)}=\|\phi\|_{W^{1/p}_q(\Gamma)}$. With this in mind,

$$\big\langle A[\gamma]\nabla(y+v)\cdot\nu,\phi\big\rangle_{W^{-1/p}_p(\Gamma)\times\mathring{W}^{1/p}_q(\Gamma)}=\int_\Omega A[\gamma]\nabla(y+v)\cdot\nabla E\phi,$$

whence

$$\big\|A[\gamma]\nabla(y+v)\cdot\nu\big\|_{W^{-1/p}_p(\Gamma)}\leq C_E C_A\,(1+\beta C_A)|v|_{W^1_p(\Omega)}\,.$$

We collect this result in the next theorem.

Theorem 4.7 (enhanced regularity). *If $G_v(u)=(\gamma,y)\in\mathbb{W}^1$ is the solution in Corollary 4.6, then $\gamma\in W^{1+1/q}_p(\Gamma)\cap\mathring{W}^1_\infty(\Gamma)$ without any further assumptions on u or v.*

4.2. G_v is Lipschitz. The first result in showing that G_v is twice Fréchet differentiable is to demonstrate that it is Lipschitz.

In the interest of saving some space we will rewrite the variational system (4.2) in the following form: find $(\gamma, y) \in \mathbb{W}^1$ such that for every $(\zeta, z) \in \mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega)$

$$\mathcal{B}_\Gamma[\gamma, \zeta] + \mathcal{B}_\Omega\left[y + v, z + E\zeta; A[\gamma]\right] = \langle u, \zeta \rangle_{W_\infty^{-1}(\Gamma) \times \mathring{W}_1^1(\Gamma)}. \tag{4.23}$$

With this new notation in place we are ready to study the Lipschitz continuity of G_v.

Theorem 4.8 (Lipschitz continuity of G_v). *If v fulfills the conditions of Corollary 4.6, then G_v satisfies*

$$\left\|G_v(u_1) - G_v(u_2)\right\|_{\mathbb{W}^1} \leq L_G \|u_1 - u_2\|_{L^2(\Gamma)} \qquad \forall u_1, u_2 \in \mathcal{U}_{ad}, \tag{4.24}$$

with constant $L_G = \frac{\alpha}{\theta_2}(1 + \beta C_A)^2 |v|_{W_p^1(\Omega)}$.

Proof. Given $u_1, u_2 \in \mathcal{U}_{ad}$, set $(\gamma_1, y_1) - (\gamma_2, y_2) = G_v(u_1) - G_v(u_2)$. Using (4.23), we have for every $(\zeta, z) \in \mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega)$

$$\begin{aligned} \mathcal{B}_\Gamma[\gamma_1 - \gamma_2, \zeta] &+ \mathcal{B}_\Omega\left[y_1 + v, z + E\zeta; A[\gamma_1]\right] \\ &- \mathcal{B}_\Omega\left[y_2 + v, z + E\zeta; A[\gamma_2]\right] = \langle u_1 - u_2, \zeta \rangle_{L^2(\Gamma) \times L^2(\Gamma)}. \end{aligned}$$

Subtracting $\mathcal{B}_\Omega\left[y_2 + v, z + E\zeta; A[\gamma_1]\right]$ from both sides and rearranging terms yields

$$\begin{aligned} \mathcal{B}_\Gamma[\gamma_1 - \gamma_2, \zeta] &+ \mathcal{B}_\Omega\left[y_1 - y_2, z + E\zeta; A[\gamma_1]\right] = \\ &+ \mathcal{B}_\Omega\left[y_2 + v, z + E\zeta; A[\gamma_2] - A[\gamma_1]\right] + \langle u_1 - u_2, \zeta \rangle_{L^2(\Gamma) \times L^2(\Gamma)}. \end{aligned}$$

The inf-sup estimates from Proposition 4.1, together with $(\gamma_i, y_i) \in \mathbb{B}_v$ for $i = 1, 2$, imply for $\zeta = 0$

$$\begin{aligned} |y_1 - y_2|_{W_p^1(\Omega)} &\leq \beta C_A |y_2 + v|_{W_p^1(\Omega)} |\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)} \\ &\leq \beta C_A (1 + \beta C_A) |v|_{W_p^1(\Omega)} |\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)}, \end{aligned} \tag{4.25}$$

and for $z = 0$

$$\begin{aligned} |\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)} &\leq \alpha C_E C_A \left(|y_1 - y_2|_{W_p^1(\Omega)} + |y_2 + v|_{W_p^1(\Omega)} |\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)}\right) \\ &\quad + \alpha \|u_1 - u_2\|_{L^2(\Gamma)} \\ &\leq \alpha C_E C_A (1 + \beta C_A)^2 |v|_{W_p^1(\Omega)} |\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)} + \alpha \|u_1 - u_2\|_{L^2(\Gamma)}. \end{aligned}$$

Finally, in view of (4.19), we infer that

$$|\gamma_1 - \gamma_2|_{W_\infty^1(\Gamma)} \leq \frac{\alpha}{\theta_2} \|u_1 - u_2\|_{L^2(\Gamma)}. \tag{4.26}$$

The asserted estimate follows immediately from the definition of $\|\cdot\|_{\mathbb{W}^1}$ in (4.11). ☐

4.3. G_v is Fréchet Differentiable. The next step towards showing the twice Fréchet differentiability of G_v entails analyzing the well-posedness of the linear variational system: find $(\gamma, y) \in \mathbb{W}^1$ such that for every $(\zeta, z) \in \mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega)$

$$\mathcal{B}_\Gamma[\gamma, \zeta] + \mathcal{D}_\Omega\left[(\gamma, y), z + E\zeta; \bar{\gamma}, \bar{y}\right] = F_\Omega(z + E\zeta) + F_\Gamma(\zeta) \tag{4.27}$$

where

$$\mathcal{D}_\Omega\left[(\gamma,y),\cdot;\bar\gamma,\bar y\right] := \mathcal{B}_\Omega\left[y,\cdot;A[\bar\gamma]\right] + \mathcal{B}_\Omega\left[\bar y + v,\cdot;\mathrm{D}A[\bar\gamma]\langle\gamma\rangle\right],$$

$(\bar\gamma,\bar y) = G_v(\bar u) \in \mathbb{B}_v$ for a fixed $\bar u$ in $\mathcal{U}_{ad}$, $\mathrm{D}A[\bar\gamma]\langle\gamma\rangle$ is given in (2.4), and $F_\Omega \in W^1_q(\Omega)^*$ and $F_\Gamma \in \mathring{W}^1_1(\Gamma)^*$ are fixed but arbitrary.

4.3.1. Preliminary Estimates. Given that the coupled system (4.27) is linear, one would be inclined to use the standard Banach-Nečas theorem to prove its well-posedness directly. We deviate from this approach and resort to the machinery already put in place.

Consider the operator $T : \mathbb{W}^1 \to \mathbb{W}^1$ given by

$$T(\gamma,y) := \big(T_1(\gamma,y), T_2(\gamma,y)\big) = (\widehat\gamma,\widehat y) \quad \forall (\gamma,y)\in\mathbb{W}^1, \tag{4.28}$$

where $\widehat\gamma = T_1(\gamma,y) \in \mathring{W}^1_\infty(\Gamma)$ satisfies for every $\zeta \in \mathring{W}^1_1(\Gamma)$

$$\mathcal{B}_\Gamma[\widehat\gamma,\zeta] = -\mathcal{D}_\Omega\left[(\gamma,y), E\zeta; A[\bar\gamma]\right] + F_\Omega(E\zeta) + F_\Gamma(\zeta), \tag{4.29}$$

and $\widehat y = T_2(\gamma,y) \in \mathring{W}^1_p(\Omega)$ satisfies for every $z \in \mathring{W}^1_q(\Omega)$

$$\mathcal{B}_\Omega\left[\widehat y, z; A[\bar\gamma]\right] = -\mathcal{B}_\Omega\left[\bar y + v, z; \mathrm{D}A[\bar\gamma]\langle T_1(\gamma,y)\rangle\right] + F_\Omega(z). \tag{4.30}$$

We point out that any fixed point of T is a also a solution to (4.27). To infer the existence of a fixed point we exploit the linear structure of (4.27). Therefore, it suffices to show the well-posedness of the intermediate operators T_1 and T_2, and to show that T is a contraction $\mathbb{W}^1$.

Lemma 4.9 (well-posedness of T_1 and T_2). *Let T_1, T_2 be the operators defined in* (4.29) *and* (4.30) *with $(\bar\gamma,\bar y)\in\mathbb{B}_v$. The following holds*

(i) for every $(\gamma,y)\in\mathbb{W}^1$, there exists a unique $\widehat\gamma = T_1(\gamma,y)$ satisfying (4.29) *and*

$$|\widehat\gamma|_{W^1_\infty(\Gamma)} \le \alpha\Big(C_E C_A\big\|(\gamma,y)\big\|_{\mathbb{W}^1} + C_E\|F_\Omega\|_{W^1_q(\Omega)^*} + \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*}\Big),$$

(ii) for every $(\gamma,y)\in\mathbb{W}^1$, there exists a unique $\widehat y = T_2(\gamma,y)$ satisfying (4.30) *and*

$$|\widehat y|_{W^1_p(\Omega)} \le \beta C_A(1+\beta C_A)|v|_{W^1_p(\Omega)}|\widehat\gamma|_{W^1_\infty(\Gamma)} + \beta\|F_\Omega\|_{W^1_q(\Omega)^*}.$$

Proof. To prove (i) we proceed as in Lemma 4.3. It suffices to check that the right hand side $\mathrm{RHS}(\zeta)$ of (4.29) is in $\mathring{W}^1_1(\Gamma)^*$, namely

$$\begin{aligned}\big|\mathrm{RHS}(\zeta)\big| \le \Bigg(& C_E C_A\Big(|y|_{W^1_p(\Omega)} + (1+\beta C_A)|v|_{W^1_p(\Omega)}|\gamma|_{W^1_\infty(\Gamma)}\Big)\\ &\qquad + C_E\|F_\Omega\|_{W^1_q(\Omega)^*} + \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*}\Bigg)|\zeta|_{W^1_1(\Gamma)}.\end{aligned}$$

The desired estimate follows from Lemma 4.2 with the coercivity of $\mathcal{B}_\Gamma$ in $\mathring{W}^1_2(\Gamma)$, and the definition of $\|\cdot\|_{\mathbb{W}^1}$ in (4.11).

Estimate (ii) is a straightforward application of the Banach-Nečas theorem [8]. □

Theorem 4.10 (T is a contraction). *Let* (4.19) *hold for some $\theta_2\in(0,1)$. The operator T defined in* (4.28) *is a contraction in $\mathbb{W}^1$ with constant $1-\theta_2$.*

Proof. We proceed in a similar fashion to Theorem 4.5. Consider not identical (γ_1, y_1) and (γ_2, y_2) in $\mathbb{W}^1$, and use (4.28) to write $(\widehat{\gamma}_i, \widehat{y}_i) = T(\gamma_i, y_i)$ for $i = 1, 2$. Applying Lemma 4.9 (i), we obtain

$$|\widehat{\gamma}_1 - \widehat{\gamma}_2|_{W^1_\infty(\Gamma)} \leq \alpha C_E C_A \big\|(\gamma_1 - \gamma_2, y_1 - y_2)\big\|_{\mathbb{W}^1} .$$

Similarly, Lemma 4.9 (ii) implies

$$|\widehat{y}_1 - \widehat{y}_2|_{W^1_p(\Omega)} \leq \beta C_A \left(1 + \beta C_A\right) |v|_{W^1_p(\Omega)} |\widehat{\gamma}_1 - \widehat{\gamma}_2|_{W^1_\infty(\Gamma)} .$$

Lastly, the upper bound (4.19) on v yields

$$\begin{aligned}\big\|(\widehat{\gamma}_1 - \widehat{\gamma}_2, \widehat{y}_1 - \widehat{y}_2)\big\|_{\mathbb{W}^1} &\leq \left(1 + \beta C_A\right)^2 |v|_{W^1_p(\Omega)} |\widehat{\gamma}_1 - \widehat{\gamma}_2|_{W^1_\infty(\Gamma)} \\ &\leq (1 - \theta_2) \big\|(\gamma_1 - \gamma_2, y_1 - y_2)\big\|_{\mathbb{W}^1} .\end{aligned}$$

Hence, T is a contraction with constant $1 - \theta_2$, as asserted. □

COROLLARY 4.11 (well-posedness of the linear system (4.27)). *Under the assumptions of Theorem 4.10, there exists a unique solution* $(\gamma, y) \in \mathbb{W}^1$ *to the variational equation* (4.27) *and the following estimates hold*

$$|\gamma|_{W^1_\infty(\Gamma)} \leq \frac{\alpha}{\theta_2} \left(C_E \left(1 + \beta C_A\right) \|F_\Omega\|_{W^1_q(\Omega)^*} + \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} \right) \tag{4.31}$$

$$|y|_{W^1_p(\Omega)} \leq \frac{\beta}{\theta_2} \left(\|F_\Omega\|_{W^1_q(\Omega)^*} + \alpha C_A \left(1 + \beta C_A\right) |v|_{W^1_p(\Omega)} \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} \right) . \tag{4.32}$$

Therefore

$$\begin{aligned}\big\|(\gamma, y)\big\|_{\mathbb{W}^1} \leq &\frac{1}{\theta_2} \left(\alpha C_E \left(1 + \beta C_A\right)^2 |v|_{W^1_p(\Omega)} + \beta \right) \|F_\Omega\|_{W^1_q(\Omega)^*} \\ &+ \frac{\alpha}{\theta_2} \left(1 + \beta C_A\right)^2 |v|_{W^1_p(\Omega)} \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} .\end{aligned}$$

Proof. Existence and uniqueness follows from Theorem 4.10. As far as the estimates, we will only derive (4.31) since the other two are merely a consequence of it. To this end we apply Lemma 4.9 and the upper bound (4.19) for v to get

$$\begin{aligned}|\gamma|_{W^1_\infty(\Gamma)} \leq\ &\alpha C_E C_A \left(\left(1 + \beta C_A\right) |v|_{W^1_p(\Omega)} |\gamma|_{W^1_\infty(\Gamma)} + |y|_{W^1_p(\Omega)} \right) \\ &+ \alpha \left(C_E \|F_\Omega\|_{W^1_q(\Omega)^*} + \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} \right) \\ \leq\ &\alpha C_E C_A \left(1 + \beta C_A\right)^2 |v|_{W^1_p(\Omega)} |\gamma|_{W^1_\infty(\Gamma)} \\ &+ \alpha C_E \left(1 + \beta C_A\right) \|F_\Omega\|_{W^1_q(\Omega)^*} + \alpha \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} . \\ \leq\ &(1 - \theta_2) |\gamma|_{W^1_\infty(\Gamma)} + \alpha C_E \left(1 + \beta C_A\right) \|F_\Omega\|_{W^1_q(\Omega)^*} + \alpha \|F_\Gamma\|_{\mathring{W}^1_1(\Gamma)^*} .\end{aligned}$$

The estimate (4.31) follows immediately. □

4.3.2. The First Order Fréchet Derivative. In this section we will prove the first order differentiability of the control to state map G_v. Since the set $\mathcal{U}_{ad}$ defined in (4.18) is not open we need to define a proper set of admissible directions.

DEFINITION 4.12. *(admissible directions) Given* $u \in \mathcal{U}_{ad}$*, the convex set* $\mathcal{C}(u)$ *comprises of all directions* $h \in L^2(\Gamma)$ *such that* $u + h \in \mathcal{U}_{ad}$*, i.e.,*

$$\mathcal{C}(u) := \left\{ h \in L^2(\Gamma) : \ u + h \in \mathcal{U}_{ad} \right\} .$$

Theorem 4.13 (the Fréchet derivative of G_v). *The control to state map $G_v : \mathcal{U}_{ad} \to \mathbb{W}^1$ is Fréchet differentiable. Its first order derivative $(\gamma, y) := G_v'(\bar{u})\, h$ in $\mathbb{W}^1$ at an arbitrary $\bar{u}$ in $\mathcal{U}_{ad}$ in the direction h in $\mathcal{C}(\bar{u})$ satisfies the linear variational system (4.27) with $F_\Omega = 0$ and $F_\Gamma(\zeta) = \int_\Gamma h\zeta$, namely*

$$\mathcal{B}_\Gamma[\gamma, \zeta] + \mathcal{D}_\Omega\big[(\gamma, y), z + E\zeta; \bar{\gamma}, \bar{y}\big] = \int_\Gamma h\zeta, \qquad \forall (\zeta, z) \in \mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega). \tag{4.33}$$

Moreover, the following estimate holds

$$\big\|(\gamma, y)\big\|_{\mathbb{W}^1} \le \frac{\alpha}{\theta_2}(1 + \beta C_A)^2 |v|_{W_p^1(\Omega)} \|h\|_{L^2(\Gamma)}\,. \tag{4.34}$$

Proof. The derivation of (4.33) is tedious but straightforward, so we skip it. The estimate (4.34) follows from Corollary 4.11.

We turn our focus to proving that G_v' is the Fréchet derivative of G_v. To this end, we must show that the remainder operator $\mathfrak{R}_{G_v}$, defined as

$$\mathfrak{R}_{G_v}[\bar{u}, h] := G_v(\bar{u} + h) - G_v(\bar{u}) - G_v'(\bar{u})\, h \qquad \forall \bar{u} \in \mathcal{U}_{ad},\ h \in \mathcal{C}(\bar{u}), \tag{4.35}$$

satisfies for all $\bar{u} \in \mathcal{U}_{ad}$

$$\lim_{h \in \mathcal{C}(\bar{u}) : \|h\|_{L^2(\Gamma)} \to 0} \frac{\big\|\mathfrak{R}_{G_v}[\bar{u}, h]\big\|_{\mathbb{W}^1}}{\|h\|_{L^2(\Gamma)}} = 0.$$

Since we do not have direct access to $\big\|\mathfrak{R}_{G_v}[\bar{u}, h]\big\|_{\mathbb{W}^1}$, the strategy of the proof is to first show that $\mathfrak{R}_{G_v}[\bar{u}, h]$ satisfies (4.27) for some $F_\Omega \in W_q^1(\Omega)^*$ and $F_\Gamma = 0$. Next, owing to the estimates in Corollary 4.11, it suffices to check that

$$\lim_{h \in \mathcal{C}(\bar{u}) : \|h\|_{L^2(\Gamma)} \to 0} \frac{\|F_\Omega\|_{W_q^1(\Omega)^*}}{\|h\|_{L^2(\Gamma)}} = 0.$$

To avoid any ambiguity we adopt the following notation in this proof,

$$\big(\gamma(\bar{u}), y(\bar{u})\big) := G_v(\bar{u}), \qquad \big(\gamma(\bar{u} + h), y(\bar{u} + h)\big) := G_v(\bar{u} + h)$$
$$\big(\gamma_u(\bar{u})h, y_u(\bar{u})h\big) := G_v'(\bar{u})h, \quad (\delta\gamma, \delta y) = \big(\mathfrak{R}_\gamma[\bar{u}, h], \mathfrak{R}_y[\bar{u}, h]\big) := \mathfrak{R}_{G_v}(\bar{u}, h),$$

whence

$$\delta\gamma = \gamma(\bar{u} + h) - \gamma(\bar{u}) - \gamma_u(\bar{u})\, h \qquad \delta y = y(\bar{u} + h) - y(\bar{u}) - y_u(\bar{u})\, h.$$

According to the remainder definition in (4.35) we start by combining (4.2) for $G_v(\bar{u} + h)$ and $G_v(\bar{u})$ with (4.33) to obtain for every (ζ, z) in $\mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega)$

$$\begin{aligned} 0 = \mathcal{B}_\Gamma\big[\gamma(\bar{u} + h) - \gamma(\bar{u}) - \gamma_u(\bar{u})h, \zeta\big] - \mathcal{D}_\Omega\Big[\big(\gamma_u(\bar{u})h, y_u(\bar{u})h\big), z + E\zeta; \gamma(\bar{u}), y(\bar{u})\Big] \\ + \mathcal{B}_\Omega\Big[y(\bar{u} + h) + v, z + E\zeta; A\big[\gamma(\bar{u} + h)\big]\Big] - \mathcal{B}_\Omega\Big[y(\bar{u}) + v, z + E\zeta; A\big[\gamma(\bar{u})\big]\Big]. \end{aligned}$$

Adding and subtracting $\mathcal{D}_\Omega\Big[\big(\gamma(\bar{u} + h) - \gamma(\bar{u}), y(\bar{u} + h) - y(\bar{u})\big), z + E\zeta; \gamma(\bar{u}), y(\bar{u})\Big]$ to the previous equation and utilizing the definition of $\delta\gamma$ and δy above, yields for every (ζ, z) in $\mathring{W}_1^1(\Gamma) \times \mathring{W}_q^1(\Omega)$

$$\mathcal{B}_\Omega[\delta\gamma, \zeta] + \mathcal{D}_\Omega\big[(\delta\gamma, \delta y), z + E\zeta; \gamma(\bar{u}), y(\bar{u})\big] = F_\Omega(z + E\zeta),$$

where

$$
\begin{aligned}
F_\Omega(\cdot) = \mathcal{B}_\Omega\Big[y(\bar u + h) + v, \cdot; A\left[\gamma(\bar u)\right] - A\left[\gamma(\bar u + h)\right]\Big] \\
+ \mathcal{B}_\Omega\Big[y(\bar u) + v, \cdot; \mathrm{D}A\left[\gamma(\bar u)\right]\left\langle \gamma(\bar u + h) - \gamma(\bar u)\right\rangle\Big].
\end{aligned}
$$

The fact that F_Ω is in $W^1_q(\Omega)^*$ follows from the continuity of $\mathcal{B}_\Omega[w, \cdot; V]$ with $|w|_{W^1_p(\Omega)}$ and $\|V\|_{L^\infty(\Omega)}$ bounded uniformly (c.f. (2.8)). Our last step is to add and subtract $\mathcal{B}_\Omega\Big[y(\bar u + h) + v, \cdot; \mathrm{D}A\left[\gamma(\bar u)\right]\left\langle \gamma(\bar u + h) - \gamma(\bar u)\right\rangle\Big]$ to F_Ω, employ the definition of the remainder $\mathfrak{R}_A$ (2.5) and the Lipschitz estimates (4.25) and (4.26) to obtain

$$
\lim_{h \in \mathcal{C}(\bar u) : \|h\|_{L^2(\Gamma)} \to 0} \frac{\Big\|\mathfrak{R}_A\left[\gamma(\bar u), \gamma(\bar u + h) - \gamma(\bar u)\right]\Big\|_{L^\infty(\Omega)}}{\|h\|_{L^2(\Gamma)}} = 0,
$$

as well as

$$
\begin{aligned}
\|F_\Omega\|_{W^1_q(\Omega)^*} &\le (1 + \beta C_A)|v|_{W^1_p(\Omega)}\Big\|\mathfrak{R}_A\left[\gamma(\bar u), \gamma(\bar u + h) - \gamma(\bar u)\right]\Big\|_{L^\infty(\Omega)} \\
&\quad + C_A\big|y(\bar u + h) - y(\bar u)\big|_{W^1_p(\Omega)}\big|\gamma(\bar u + h) - \gamma(\bar u)\big|_{W^1_\infty(\Gamma)} \\
&= o\Big(\|h\|_{L^2(\Gamma)}\Big).
\end{aligned}
$$

This concludes the proof. ☐

4.4. The Second Order Fréchet Derivative. The final result of this section is to show that $G_v(u)$ is twice Fréchet differentiable with respect to u. The technique is similar to the one in subsections 4.2 and 4.3. In fact, proceeding as in Theorem 4.8 we get the following.

Proposition 4.14 (Lipschitz continuity of G'_v). *There exists a constant $L_{G'} > 0$, such that for every $u_1, u_2 \in \mathcal{U}_{ad}$*

$$
\sup_{0 \ne h \in \mathcal{C}(u_1) \cap \mathcal{C}(u_2)} \frac{\left\|G'_v(u_1)h - G'_v(u_2)h\right\|_{\mathbb{W}^1}}{\|h\|_{L^2(\Gamma)}} \le L_{G'}\|u_1 - u_2\|_{L^2(\Gamma)}. \tag{4.36}
$$

Theorem 4.15 (the Fréchet derivative of $G'_v(\bar u)h$). *The control to state map $G_v : \mathcal{U}_{ad} \to \mathbb{W}^1$ is twice Fréchet differentiable. Its second order derivative $(\gamma, y) := G''_v(\bar u)h_1h_2$ in $\mathbb{W}^1$ at an arbitrary $\bar u \in \mathcal{U}_{ad}$ in the direction $(h_1, h_2) \in \mathcal{C}(\bar u) \times \mathcal{C}(\bar u)$ satisfies the linear variational system (4.27), namely for every (ζ, z) in $\mathring{W}^1_1(\Gamma) \times \mathring{W}^1_q(\Omega)$*

$$
\mathcal{B}_\Gamma[\gamma, \zeta] + \mathcal{D}_\Omega\left[(\gamma, y), z + E\zeta; \bar\gamma, \bar y\right] = F_\Omega(z + E\zeta), \tag{4.37}
$$

with $F_\Omega \in W^1_q(\Omega)^*$ given by

$$
\begin{aligned}
F_\Omega(\cdot) := -\mathcal{B}_\Omega\left[y_1, \cdot; \mathrm{D}A[\bar\gamma]\langle\gamma_2\rangle\right] \\
- \mathcal{B}_\Omega\left[y_2, \cdot; \mathrm{D}A[\bar\gamma]\langle\gamma_1\rangle\right] - \mathcal{B}_\Omega\Big[\bar y + v, \cdot; \mathrm{D}^2A[\bar\gamma]\langle\gamma_1, \gamma_2\rangle\Big],
\end{aligned} \tag{4.38}
$$

and $(\gamma_i, y_i) := G'_v(\bar u)h_i$, for $i = 1, 2$. Moreover, the following estimates hold

$$
|\gamma|_{W^1_\infty(\Gamma)} \le \frac{\alpha^3}{\theta_2^3}C_EC_A(1 + 2\beta C_A)(1 + \beta C_A)^2|v|_{W^1_p(\Omega)}\|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)}, \tag{4.39}
$$

$$
|y|_{W^1_p(\Omega)} \le \frac{\alpha^2}{\theta_2^3}\beta C_A(1 + 2\beta C_A)(1 + \beta C_A)|v|_{W^1_p(\Omega)}\|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)}. \tag{4.40}
$$

Proof. We skip the derivation of (4.37) because it is tedious but straightforward. The estimates for $|\gamma|_{W^1_\infty(\Gamma)}$ and $|y|_{W^1_p(\Omega)}$ are a consequence of Corollary 4.11 with $F_\Gamma = 0$ after estimating (4.38), namely

$$
\begin{aligned}
\|F_\Omega\|_{W^1_q(\Omega)^*} &\leq C_A \left(|y_1|_{W^1_p(\Omega)}|\gamma_2|_{W^1_\infty(\Gamma)} + |y_2|_{W^1_p(\Omega)}|\gamma_1|_{W^1_\infty(\Gamma)}\right) \\
&\quad + C_A\left(1+\beta C_A\right)|v|_{W^1_p(\Omega)}|\gamma_1|_{W^1_\infty(\Gamma)}|\gamma_2|_{W^1_\infty(\Gamma)} \\
&\leq 2\left(\frac{\alpha}{\theta_2}\right)^2 \beta C_A^2\left(1+\beta C_A\right)|v|_{W^1_p(\Omega)}\|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)} \\
&\quad + \left(\frac{\alpha}{\theta_2}\right)^2 C_A\left(1+\beta C_A\right)|v|_{W^1_p(\Omega)}\|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)} \\
&= \left(\frac{\alpha}{\theta_2}\right)^2 C_A\left(1+2\beta C_A\right)\left(1+\beta C_A\right)|v|_{W^1_p(\Omega)}\|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)},
\end{aligned}
$$

where we have used (4.31)-(4.32) with $F_\Omega = 0$ for (γ_i, y_i) along with (4.6).

The strategy for showing second order Frechét differentiability of G_v is the same as in Theorem 4.13: we first show that the remainder

$$
(\delta\gamma, \delta y) := G'_v\left(\bar{u}+h_2\right)h_1 - G'_v\left(\bar{u}\right)h_1 - G''_v\left(\bar{u}\right)h_1h_2, \tag{4.41}
$$

satisfies the linear variational system in (4.27) for a suitable right-hand side $\delta F_\Omega \in W^1_q\left(\Omega\right)^*$, and prove that

$$
\sup_{0\neq h_1\in\mathcal{C}(\bar{u})} \frac{\|\delta F_\Omega\|_{W^1_q(\Omega)^*}}{\|h_1\|_{L^2(\Gamma)}} = o\left(\|h_2\|_{L^2(\Gamma)}\right). \tag{4.42}
$$

As a tradeoff between clarity and space we denote $u_0 = \bar{u}$, $u_i = \bar{u} + h_i$, and

$$
\begin{aligned}
\big(\gamma(u_i), y(u_i)\big) &:= G_v(u_i), \\
\big(\gamma_u\left(u_i\right)h_j, y_u\left(u_i\right)h_j\big) &:= G'_v\left(u_i\right)h_j \\
\big(\gamma_{uu}\left(u_0\right)h_1h_2, y_{uu}\left(u_0\right)h_1h_2\big) &:= G''_v\left(u_0\right)h_1h_2,
\end{aligned}
$$

for $i, j = 1, 2$, whence

$$
\begin{aligned}
\delta\gamma &= \gamma_u\left(u_2\right)h_1 - \gamma_u\left(u_0\right)h_1 - \gamma_{uu}\left(u_0\right)h_1h_2, \\
\delta y &= y_u\left(u_2\right)h_1 - y_u\left(u_0\right)h_1 - y_{uu}\left(u_0\right)h_1h_2.
\end{aligned}
$$

According to the remainder definition in (4.41) we start by combining (4.33) for $G'_v\left(u_2\right)h_1$ and $G'_v\left(u_0\right)h_1$ with (4.37) to obtain for every (ζ, z) in $\mathring{W}^1_1\left(\Gamma\right)\times\mathring{W}^1_q\left(\Omega\right)$

$$
\begin{aligned}
-F_\Omega\left(z+E\zeta\right) = \mathcal{B}_\Gamma\left[\delta\gamma,\zeta\right] &+ \mathcal{D}_\Omega\left[G'_v(u_2)h_1, z+E\zeta; G_v(u_2)\right] \\
&- \mathcal{D}_\Omega\left[G'_v(u_0)h_1 + G''_v(u_0)h_1h_2, z+E\zeta; G_v(u_0)\right],
\end{aligned}
$$

where $-F_\Omega(\cdot) = \sum_{i=1}^3 F_i(\cdot)$ is defined in (4.38). Further manipulation, based on adding to both sides the following two additional terms,

$$
\begin{aligned}
F_4\left(z+E\zeta\right) &= \mathcal{D}_\Omega\left[G'_v(u_2)h_1, z+E\zeta; G_v(u_0)\right], \\
F_5\left(z+E\zeta\right) &= -\mathcal{D}_\Omega\left[G'_v(u_2)h_1, z+E\zeta; G_v(u_2)\right]
\end{aligned}
$$

leads to

$$\mathcal{B}_\Gamma\left[\delta\gamma,\zeta\right]+\mathcal{D}_\Omega\left[\left(\delta\gamma,\delta y\right),z+E\zeta;\gamma(u_0),y(u_0)\right]=-\delta F_\Omega\left(z+E\zeta\right),$$

where $\delta F_\Omega=\sum_{i=1}^{5}F_i$ is clearly in $W_q^1\left(\Omega\right)^*$. To create additional cancellations we further decompose $\delta F_\Omega=\sum_{i=1}^{9}T_i$ as follows:

$$\begin{aligned}
T_1&=\mathcal{B}_\Omega\Big[y_u(u_0)h_1,\cdot;\mathrm{D}A\left[\gamma(u_0)\right]\left\langle\gamma(u_2)-\gamma(u_0)-\gamma_u(u_0)h_2\right\rangle\Big],\\
T_2&=\mathcal{B}_\Omega\Big[y_u(u_2)h_1-y_u(u_0)h_1,\cdot;\mathrm{D}A\left[\gamma(u_0)\right]\left\langle\gamma(u_2)-\gamma(u_0)\right\rangle\Big],\\
T_3&=\mathcal{B}_\Omega\Big[y_u(u_2)h_1,\cdot;A\left[\gamma(u_2)\right]-A\left[\gamma(u_0)\right]-\mathrm{D}A\left[\gamma(u_0)\right]\left\langle\gamma(u_2)-\gamma(u_0)\right\rangle\Big],\\
T_4&=\mathcal{B}_\Omega\Big[y_u(u_0)h_2,\cdot;\mathrm{D}A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_2)h_1-\gamma_u(u_0)h_1\right\rangle\Big],\\
T_5&=\mathcal{B}_\Omega\Big[y(u_2)-y(u_0)-y_u(u_0)h_2,\cdot;\mathrm{D}A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_2)h_1\right\rangle\Big],\\
T_6&=\mathcal{B}_\Omega\Big[y(u_2)+v,\cdot;\Big(\mathrm{D}A\left[\gamma(u_2)\right]-\mathrm{D}A\left[\gamma(u_0)\right]\Big)\left\langle\gamma_u(u_2)h_1\right\rangle\\
&\qquad\qquad-\mathrm{D}^2A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_2)h_1,\gamma(u_2)-\gamma(u_0)\right\rangle\Big],\\
T_7&=\mathcal{B}_\Omega\left[y(u_2)+v,\cdot;\mathrm{D}^2A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_2)h_1,\gamma(u_2)-\gamma(u_0)-\gamma_u(u_0)h_2\right\rangle\Big]\right.,\\
T_8&=\mathcal{B}_\Omega\Big[y(u_2)+v,\cdot;\mathrm{D}^2A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_2)h_1-\gamma_u(u_0)h_1,\gamma_u(u_0)h_2\right\rangle\Big],\\
T_9&=\mathcal{B}_\Omega\Big[y(u_2)-y(u_0),\cdot;\mathrm{D}^2A\left[\gamma(u_0)\right]\left\langle\gamma_u(u_0)h_1,\gamma_u(u_0)h_2\right\rangle\Big].
\end{aligned}$$

We now estimate each of these terms separately and show

$$\sup_{0\neq h_1\in\mathcal{C}(u_0)}\frac{\|T_i\|_{W_q^1(\Omega)^*}}{\|h_1\|_{L^2(\Gamma)}}=o\Big(\|h_2\|_{L^2(\Gamma)}\Big). \tag{4.43}$$

which obviously imply (4.42).

- Term T_1: Since

  $$\|T_1\|_{W_q^1(\Omega)^*}\le C_A\big|y_u\left(u_0\right)h_1\big|_{W_p^1(\Omega)}\big|\gamma\left(u_2\right)-\gamma\left(u_0\right)-\gamma_u\left(u_0\right)h_2\big|_{W_\infty^1(\Gamma)},$$

  the estimate (4.34), together with

  $$\big|\gamma\left(u_2\right)-\gamma\left(u_0\right)-\gamma_u\left(u_0\right)h_2\big|_{W_\infty^1(\Gamma)}=o\Big(\|h_2\|_{L^2(\Gamma)}\Big),$$

  implies (4.43).
- Term T_2: Since

  $$\|T_2\|_{W_q^1(\Omega)^*}\le C_A\big|y_u\left(u_2\right)h_1-y_u\left(u_0\right)h_1\big|_{W_p^1(\Omega)}\big|\gamma\left(u_2\right)-\gamma\left(u_0\right)\big|_{W_\infty^1(\Gamma)},$$

  it suffices to recall the Lipschitz properties (4.24) of G_v, and (4.36) of G_v' to deduce (4.43).
- Term T_3: Invoking the Fréchet differentiability (2.5) of A, and the Lipschitz property (4.26) of $\gamma(u_0)$ we infer that

  $$\begin{aligned}\Big\|A\left[\gamma\left(u_2\right)\right]-A\left[\gamma\left(u_0\right)\right]-\mathrm{D}A\left[\gamma\left(u_0\right)\right]\left\langle\gamma\left(u_2\right)-\gamma\left(u_0\right)\right\rangle\Big\|_{L^\infty(\Omega)}&\\
  =o\Big(\big|\gamma\left(u_2\right)-\gamma\left(u_0\right)\big|_{W_\infty^1(\Gamma)}\Big)=o\Big(\|h_2\|_{L^2(\Gamma)}\Big).&\end{aligned}$$

This, in conjuction with $\left|y_u\left(u_2\right)h_1\right|_{W^1_p(\Omega)} \lesssim \|h_1\|_{L^2(\Gamma)}$, yields (4.43).

- Term T_4: In view of the Lipschitz property (4.36) of G'_v
$$\left|\gamma_u\left(u_2\right)h_1 - \gamma_u\left(u_0\right)h_1\right|_{W^1_\infty(\Gamma)} \lesssim \|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)}\,,$$
property (4.43) follows from $\left|y_u\left(u_0\right)h_2\right|_{W^1_p(\Omega)} \lesssim \|h_2\|_{L^2(\Gamma)}$.
- Term T_5: Since $y(u)$ is Fréchet differentiable according to Theorem 4.13, namely
$$\left|y\left(u_2\right) - y\left(u_0\right) - y_u\left(u_0\right)h_2\right|_{W^1_p(\Omega)} = o\left(\|h_2\|_{L^2(\Gamma)}\right),$$
the bound (4.43) is a consequence of $\left|\gamma_u\left(u_2\right)h_1\right|_{W^1_\infty(\Gamma)} \lesssim \|h_1\|_{L^2(\Gamma)}$.
- Term T_6: We recall the second order Fréchet differentiability (2.7) of A , and the Lipschitz continuity (4.24) of G_v, to write
$$\begin{aligned}&\Big\| \mathrm{D}A\left[\gamma\left(u_2\right)\right]\langle\gamma_u(u_2)h_1\rangle - \mathrm{D}A\left[\gamma\left(u_0\right)\right]\langle\gamma_u(u_2)h_1\rangle \\ &\qquad - \mathrm{D}^2A\left[\gamma\left(u_0\right)\right]\langle\gamma_u\left(u_2\right)h_1, \gamma\left(u_2\right) - \gamma\left(u_0\right)\rangle\Big\|_{L^\infty(\Gamma)} \\ &\qquad = \|h_1\|_{L^2(\Gamma)}\, o\left(\left|\gamma\left(u_2\right) - \gamma\left(u_0\right)\right|_{W^1_\infty(\Gamma)}\right) = \|h_1\|_{L^2(\Gamma)}\, o\left(\|h_2\|_{L^2(\Gamma)}\right).\end{aligned}$$
Since $\left|y(u_2) + v\right|_{W^1_p(\Omega)} \lesssim |v|_{W^1_p(\Omega)}$, this implies (4.43).
- Term T_7: We proceed as with T_6, now appealing to (2.8) and the Fréchet differentiability of γ at u_0 (Theorem 4.13), to obtain
$$\left|\gamma_u(u_2)h_1\right|_{W^1_\infty(\Gamma)}\left|\gamma(u_2) - \gamma(u_0) - \gamma_u(u_0)h_2\right|_{W^1_\infty(\Gamma)} = \|h_1\|_{L^2(\Gamma)}\, o\left(\|h_2\|_{L^2(\Gamma)}\right),$$
whence (4.43).
- Term T_8: We employ the Lipschitz property (4.36) of G'_v in to write
$$\left|\gamma_u(u_2)h_1 - \gamma_u(u_0)h_1\right|_{W^1_\infty(\Gamma)} \lesssim \|h_1\|_{L^2(\Gamma)}\|h_2\|_{L^2(\Gamma)}\,.$$
The desired bound (4.43) follows from $\left|\gamma_u\left(u_0\right)h_2\right|_{W^1_\infty(\Gamma)} \lesssim \|h_2\|_{L^2(\Gamma)}$.
- Term T_9: We use the Lipschitz property (4.24) of G_v,
$$\left|y(u_2) - y(u_0)\right|_{W^1_p(\Omega)} \lesssim \|h_2\|_{L^2(\Gamma)}\,,$$
together with $\left|\gamma_u\left(u_0\right)h_j\right|_{W^1_\infty(\Gamma)} \lesssim \left\|h_j\right\|_{L^2(\Gamma)}$ to deduce (4.43).

Altogether, this concludes the proof. ☐

5. Optimal Control. Let us summarize what we have accomplished so far. We have formally derived the first order necessary optimality conditions in section 3. If G_v denotes the control to state map, we have proved in section 4 that G_v is well posed, i.e., there exists a unique weak solution to the state equations (2.3b) for every $u \in \mathcal{U}_{ad}$ in (4.18), and v satisfying (4.16) and (4.19). As a crucial step forward we have shown that G_v is twice Fréchet differentiable.

This background work puts us in the position to show the existence and uniqueness of the optimal control u solving the OC-FBP in (2.3a)-(2.3b). We will achieve this result in three stages. We first show the existence of u in Theorem 5.1 of subsection 5.1. We next derive the first order necessary optimality conditions and show the existence of solution to the adjoint equations in subsection 5.2. Finally in subsection 5.3 we end this voyage by proving the second order sufficient conditions for the control u.

5.1. Existence of Optimal Control. In order to show the existence of a solution to our optimal control problem we must first rewrite the cost functional $\mathcal{J} : \mathbb{W}^1 \times \mathcal{U}_{ad} \to \mathbb{R}$ from (2.3a) in its reduced form. This is accomplished by utilizing the control to state map G_v from Section 4 as follows:

$$\mathcal{J}(u) := \mathcal{J}(G_v(u), u) = \mathcal{J}(\gamma, y, u) = \mathcal{J}_1(G_v(u)) + \mathcal{J}_2(u),$$

with

$$\mathcal{J}_1(G_v(u)) := \frac{1}{2}\|\gamma - \gamma_d\|^2_{L^2(\Gamma)} + \frac{\lambda_y}{2}\|y + v - y_d\|^2_{L^2(\Omega)}, \quad \mathcal{J}_2(u) := \frac{\lambda_u}{2}\|u\|^2_{L^2(\Gamma)}.$$

Now an equivalent minimization problem to (2.3a) is

$$\min_{u \in \mathcal{U}_{ad}} \mathcal{J}(u). \tag{5.1}$$

THEOREM 5.1. *For every v satisfying* (4.16), (4.19), *there exists an optimal control $\bar{u} \in \mathcal{U}_{ad}$ minimizing the cost functional* (2.3a) *with optimal state $(\bar{\gamma}, \bar{y}) \in \left(W^{1+1/q}_p(\Gamma) \cap \mathring{W}^1_\infty(\Gamma)\right) \times \mathring{W}^1_p(\Omega)$ which solves the free boundary problem* (2.3b) *and satisfies the state constraint* (1.2c).

Proof. In order to show the existence of an optimal control we use the direct method of the calculus of variations. We first note that the cost functional $\mathcal{J}$ in (5.1) is bounded below by zero, whence $j = \inf_{u \in \mathcal{U}_{ad}} \mathcal{J}(u)$ is finite. We thus construct a minimizing sequence $\{u_n\}_{n \in \mathbb{N}}$ such that

$$j = \lim_{n \to \infty} \mathcal{J}(u_n).$$

By Definition 4.4, $\mathcal{U}_{ad}$ is nonempty, closed, bounded and convex in $L^2(\Gamma)$, thus weakly sequentially compact. Consequently, we can extract a weakly convergent subsequence $\{u_{n_k}\}_{k \in \mathbb{N}} \subset L^2(\Gamma)$, i.e.

$$u_{n_k} \rightharpoonup \bar{u} \quad \text{in } L^2(\Gamma), \qquad \bar{u} \in \mathcal{U}_{ad}.$$

Here $\bar{u}$ is our optimal control candidate.

Henceforth, we drop the subindex k when extracting subsequences. According to Corollary 4.6 and (4.22), we let $G_v(u_n) = (\gamma_n, y_n) \in \left(W^{1+1/q}_p(\Gamma) \cap \mathring{W}^1_\infty(\Gamma)\right) \times \mathring{W}^1_p(\Omega)$ denote the unique state corresponding to u_n, thereby solving the free boundary problem (2.3b) and satisfying the state constraint (1.2c). Since $W^{1+1/q}_p(\Gamma) \cap \mathring{W}^1_\infty(\Gamma)$ is compactly embedded into $\mathring{W}^1_\infty(\Gamma)$, using Rellich-Kondrachov theorem we can extract a strongly convergent subsequence $\{\gamma_n\}_{n \in \mathbb{N}} \subset \mathring{W}^1_\infty(\Gamma)$, i.e.

$$\gamma_n \to \bar{\gamma} \quad \text{in } \mathring{W}^1_\infty(\Gamma), \qquad \text{and} \qquad y_n \rightharpoonup \bar{y} \quad \text{in } \mathring{W}^1_p(\Omega).$$

Note that the limit pair $(\bar{\gamma}, \bar{y})$ is the state corresponding to the control $\bar{u}$. This results from replacing (γ, y) with (γ_n, y_n) in the variational equation (4.2) taking the limit, and making use of the embedding $L^2(\Gamma) \subset W^{-1}_\infty(\Gamma)$.

Finally, using the fact that $\mathcal{J}_2(u)$ is continuous in L^2 and convex, together with the the strong convergence $(\gamma_n, y_n) \to (\bar{\gamma}, \bar{y})$ in $L^2(\Gamma) \times L^2(\Omega)$ due to Rellich-Kondrachov theorem, it follows that $\mathcal{J}$ is weakly lower semicontinuous, whence

$$\inf_{u \in \mathcal{U}_{ad}} \mathcal{J}(u) = \liminf_{n \to \infty} \big(\mathcal{J}_1(G_v(u_n)) + \mathcal{J}_2(u_n)\big) \geq \mathcal{J}_1(G_v(\bar{u})) + \mathcal{J}_2(\bar{u}) = \mathcal{J}(\bar{u}).$$

This concludes the proof. □

5.2. First Order Necessary Condition.

We start with a classical result [19].

Lemma 5.2 (variational inequality). *If $\bar{u} \in \mathcal{U}_{ad}$ denotes an optimal control, given by Theorem 5.1 then the first order necessary optimality condition satisfied by $\bar{u}$ is*

$$\left\langle \mathcal{J}'(\bar{u}), u - \bar{u} \right\rangle_{L^2(\Gamma)\times L^2(\Gamma)} \geq 0 \quad \forall u \in \mathcal{U}_{ad}. \tag{5.2}$$

We will show that the variational inequality (5.2) is the same as (3.7) and in the process we will prove that (3.5) and (3.6) are the correct adjoint equations, thereby furnishing a rigorous derivation of the formal results in Section 3.

Theorem 5.3 (first order conditions). *If $\bar{u} \in \mathcal{U}_{ad}$ denotes an optimal control of OC-FBP, then the first order necessary optimality conditions are given by* (3.5), (3.6) *and* (3.7).

Proof. We can infer that $\mathcal{J}$ is Fréchet differentiable by recalling from Theorems 4.13 and 4.15 that G_v is twice differentiable and that $\mathcal{J}_1$ is quadratic. In fact, the Fréchet derivative of $\mathcal{J}$ in (5.1) at $\bar{u}$ in a direction h is

$$\mathcal{J}'(\bar{u})h = \mathcal{J}_1'(G_v(\bar{u}))G_v'(\bar{u})h + \mathcal{J}_2'(\bar{u})h = \mathcal{J}_1'(G_v(\bar{u}))(\gamma_u, y_u) + \mathcal{J}_2'(\bar{u})h,$$

where $(\gamma_u, y_u) = G_v'(\bar{u})h$ satisfies (4.33) and

$$\begin{aligned} \mathcal{J}'(\bar{u})h = {} & \lambda_y \langle \bar{y} + v - y_d, y_u \rangle_{L^2(\Omega)\times L^2(\Omega)} \\ & + \langle \bar{\gamma} - \gamma_d, \gamma_u \rangle_{L^2(\Gamma)\times L^2(\Gamma)} + \lambda_u \langle \bar{u}, h \rangle_{L^2(\Gamma)\times L^2(\Gamma)}. \end{aligned} \tag{5.3}$$

Introducing the adjoint states $(\bar{r}, \bar{s}) \in W_q^1(\Omega) \times \mathring{W}_1^1(\Gamma)$, which satisfy the system (3.5)-(3.6) in weak form, and noting that $h \in L^2(\Gamma)$, we obtain

$$\mathcal{J}'(\bar{u})h = \mathcal{B}_\Gamma\left[\gamma_u, \bar{s}\right] + \mathcal{D}_\Omega\left[\left(\gamma_u, y_u\right), \bar{r}; \bar{\gamma}, \bar{y}\right] + \lambda_u \langle \bar{u}, h \rangle_{L^2(\Gamma)\times L^2(\Gamma)}.$$

Utilizing (4.33) with $\zeta = \bar{s}$ and $z = \bar{r}$, we arrive at

$$\mathcal{J}'(\bar{u})h = \langle \bar{s} + \lambda_u \bar{u}, h \rangle_{L^2(\Gamma)\times L^2(\Gamma)} + \mathcal{D}_\Omega\left[\left(\gamma_u, y_u\right), \bar{r} - E\bar{s}; \bar{\gamma}, \bar{y}\right].$$

Since the Dirichlet condition $\bar{r}|_\Gamma = \bar{s}$ implies $\bar{r} - E\bar{s} \in \mathring{W}_q^1(\Omega)$, (4.33) with $\zeta = 0$ and $z \in \mathring{W}_q^1(\Omega)$ yields $\mathcal{D}_\Omega\left[\left(\gamma_u, y_u\right), z; \bar{\gamma}, \bar{y}\right] = 0$, whence

$$\mathcal{J}'(\bar{u})h = \langle \bar{s} + \lambda_u \bar{u}, h \rangle_{L^2(\Gamma)\times L^2(\Gamma)}.$$

In view of (5.2), this coincides with (3.7) for $h = u - \bar{u}$ admissible. □

5.2.1. Well-posedness of the Adjoint System.

Before we dwell upon the second order sufficient optimality conditions we put together the last piece of the puzzle: the well-posedness of the adjoint system (3.5) and (3.6). This will be done using a contraction argument in Banach spaces, assuming that we have a solution $(\bar{\gamma}, \bar{y}) \in \mathbb{B}_v$ to the state equations in (2.3b) satisfying Proposition 4.1 and Lemma 4.3.

Let $\mathbb{V} := \left\{ r \in W_q^1(\Omega) : r|_\Gamma \in \mathring{W}_1^1(\Gamma),\ r|_\Sigma = 0 \right\}$, and the operator $T_1 : \mathbb{V} \to \mathring{W}_1^1(\Gamma)$ be defined as $\widetilde{s} = T_1(r)$ where $\widetilde{s}$ satisfies for every $\zeta \in \mathring{W}_\infty^1(\Gamma)$

$$\mathcal{B}_\Gamma\left[\zeta, \widetilde{s}\right] = \left\langle \zeta, \mathbf{f}\left[\bar{\gamma}, \bar{y}, r\right] \right\rangle_{\mathring{W}_\infty^1(\Gamma), W_1^{-1}(\Gamma)}, \tag{5.4}$$

with

$$\langle \zeta, \mathbf{f} \rangle_{\mathring{W}^1_\infty(\Gamma), W_1^{-1}(\Gamma)} := \left\langle \zeta, f_0\left[x_1; \bar{\gamma}, \bar{y}, r\right] \right\rangle_{L^\infty(\Gamma), L^1(\Gamma)} + \left\langle \mathrm{d}_{x_1} \zeta, f_1\left[x_1; \bar{\gamma}, \bar{y}, r\right] \right\rangle_{L^\infty(\Gamma), L^1(\Gamma)}$$

and

$$\begin{aligned} f_0[\cdot; \bar{\gamma}, \bar{y}, r] &:= \bar{\gamma} - \gamma_d - \int_0^1 A_1\left[\bar{\gamma}\right] \nabla \bar{y} \cdot \nabla r \, \mathrm{d}x_2, \\ f_1[\cdot; \bar{\gamma}, \bar{y}, r] &:= -\int_0^1 A_2\left[\bar{\gamma}\right] \nabla \bar{y} \cdot \nabla r \, \mathrm{d}x_2. \end{aligned} \tag{5.5}$$

Given $\widetilde{s} \in \mathring{W}_1^1(\Gamma)$, let $T_2 : \mathring{W}_1^1(\Gamma) \to \mathbb{V}$ be the operator defined as $\widetilde{r} = T_2(\widetilde{s}) = \ell + E\widetilde{s}$ with $\ell \in \mathring{W}_q^1(\Omega)$ satisfying

$$\mathcal{B}_\Omega\left[z, \widetilde{r}; A\left[\bar{\gamma}\right]\right] = \lambda_y \langle z, \bar{y} + v - y_d \rangle_{L^p(\Omega) \times L^q(\Omega)} \qquad \forall z \in \mathring{W}_p^1(\Omega). \tag{5.6}$$

Lemma 5.4 (ranges of T_1 and T_2). *Let T_1, T_2 be defined in* (5.4) *and* (5.6). *If $(\bar{\gamma}, \bar{y}) \in \mathbb{B}_v$, defined in* (4.12), *then for every*

(i) $r \in \mathbb{V}$, the solution $\widetilde{s} = T_1(r)$ to (5.4) *satisfies*

$$|\widetilde{s}|_{W_1^1(\Gamma)} \le \alpha\left(\|\bar{\gamma} - \gamma_d\|_{L^1(\Gamma)} + 2\beta C_A^2 |v|_{W_p^1(\Omega)} |r|_{W_q^1(\Omega)}\right);$$

(ii) $\widetilde{s} \in \mathring{W}_1^1(\Gamma)$, the solution $\widetilde{r} = T_2(\widetilde{s})$ to (5.6) *satisfies*

$$|\widetilde{r}|_{W_q^1(\Omega)} \le \beta\left(\|\bar{y} + v - y_d\|_{L^q(\Omega)} + C_E C_A |\widetilde{s}|_{W_1^1(\Gamma)}\right).$$

Proof. Using (4.8) of Proposition 4.1, and applying Banach-Nečas theorem [8], there exists a unique solution $\widetilde{r}$ to (5.6). Estimate (ii) follows from (4.8) and (2.8), as well as the Poincaré inequality $\|z\|_{L^p(\Omega)} \le |z|_{W_p^1(\Omega)}$ for the unit square.

In order to show the existence of solution to (5.4) we note that we are looking for an absolutely continuous function on $\Gamma = (0,1)$ with zero Dirichlet values. Therefore, by the characterization of such functions in $\mathbb{R}$ [15, Theorem 5.14], there exists a $g \in L^1(0,1)$ such that

$$\widetilde{s}(x_1) = \int_0^{x_1} \left(g(t) - \int_0^1 g(\tau)\, \mathrm{d}\tau\right) \mathrm{d}t \qquad \forall x_1 \in (0,1)$$

because $s_1(0) = s_1(1) = 0$. The variational equation (5.4) is satisfied by $\widetilde{s}$ with

$$g(t) = -\frac{1}{\kappa}\left(f_1[t; \bar{\gamma}, \bar{y}, r] + \int_0^t f_0[\tau; \bar{\gamma}, \bar{y}, r]\, \mathrm{d}\tau\right)$$

and f_0, f_1 defined in (5.5).

It remains to check that $\int_0^t f_0 d\tau$ and f_1 are in $L^1(0,1)$. This follows by applying Fubini's theorem and Hölder's inequality. We consider first f_0:

$$\begin{aligned} \int_0^1 \left|f_0[x_1; \cdot]\right| \mathrm{d}x_1 &\le \|\bar{\gamma} - \gamma_d\|_{L^1(0,1)} + \int_\Omega \left|A_1\left[\bar{\gamma}\right] \nabla \bar{y} \cdot \nabla r\right| \mathrm{d}x \\ &\le \|\bar{\gamma} - \gamma_d\|_{L^1(0,1)} + C_A |\bar{y}|_{W_p^1(\Omega)} |r|_{W_q^1(\Omega)} \\ &< \|\bar{\gamma} - \gamma_d\|_{L^1(0,1)} + \beta C_A^2 |v|_{W_p^1(\Omega)} |r|_{W_q^1(\Omega)}, \end{aligned}$$

because $|\bar{y}|_{W^1_p(\Omega)} \le \beta C_A |v|_{W^1_p(\Omega)}$. Similarly, we obtain an L^1 estimate for f_1

$$\int_0^1 \big|f_1(x_1)\big|\,\mathrm{d}x_1 \le \int_\Omega \big|A_2\left[\bar{\gamma}\right] \nabla \bar{y} \cdot \nabla r\big|\,\mathrm{d}x \le \beta C_A^2 |v|_{W^1_p(\Omega)} |r|_{W^1_q(\Omega)}\,.$$

This implies (i). Finally, the uniqueness of $\widetilde{s}$ follows from the estimate in (4.7b). □

THEOREM 5.5 (existence of (3.5) and (3.6)). *Under the assumptions of Lemma 5.4, the operator* $T = T_2 \circ T_1 : \mathbb{V} \to \mathbb{V}$ *is a contraction with constant* $\beta C_A/\left(1+\beta C_A\right)$.

Proof. The proof is similar to the one in Theorem 4.5, therefore we will be brief. Consider r_1, $r_2 \in \mathbb{V}$ such that $r_1 \neq r_2$ and let $\widetilde{s}_i = T_1(r_i), \widetilde{r}_i = T_2(\widetilde{s}_i)$, where $\widetilde{s}_i, \widetilde{r}_i$ solve (5.4) and (5.6) for $i = 1, 2$. Then Proposition 4.1 (i) and Lemma 5.4 (i) imply

$$\begin{aligned}
|\widetilde{s}_1 - \widetilde{s}_2|_{W^1_1(\Gamma)} &\le \alpha \sup_{|\zeta|_{W^1_\infty(\Gamma)}=1} \mathcal{B}_\Gamma\left[\zeta, \widetilde{s}_1 - \widetilde{s}_2\right] \\
&= \alpha \sup_{|\zeta|_{W^1_\infty(\Gamma)}=1} \big\langle \zeta, \mathbf{f}\left[\bar{\gamma}, \bar{y}, r_1\right] - \mathbf{f}\left[\bar{\gamma}, \bar{y}, r_2\right] \big\rangle \\
&\le \alpha \beta C_A^2 |v|_{W^1_p(\Omega)} |r_1 - r_2|_{W^1_q(\Omega)}\,.
\end{aligned}$$

In addition, since

$$|\widetilde{r}_1 - \widetilde{r}_2|_{W^1_q(\Omega)} = \Big|\tilde{\ell}_1 + E\widetilde{s}_1 - \tilde{\ell}_2 - E\widetilde{s}_2\Big|_{W^1_q(\Omega)} \le C_E |\widetilde{s}_1 - \widetilde{s}_2|_{W^1_1(\Gamma)} + \Big|\tilde{\ell}_1 - \tilde{\ell}_2\Big|_{W^1_q(\Omega)}\,,$$

and Proposition 4.1 (ii) implies

$$\begin{aligned}
\Big|\tilde{\ell}_1 - \tilde{\ell}_2\Big|_{W^1_q(\Omega)} &\le \beta \sup_{|z|_{W^1_p(\Omega)}=1} \mathcal{B}_\Omega\left[z, \tilde{\ell}_1 - \tilde{\ell}_2; A\left[\bar{\gamma}\right]\right] \\
&= \beta \sup_{|z|_{W^1_p(\Omega)}=1} \mathcal{B}_\Omega\left[z, E\widetilde{s}_2 - E\widetilde{s}_1; A\left[\bar{\gamma}\right]\right],
\end{aligned}$$

we deduce that

$$\begin{aligned}
|\widetilde{r}_1 - \widetilde{r}_2|_{W^1_q(\Omega)} &\le C_E\left(1 + \beta C_A\right) |\widetilde{s}_1 - \widetilde{s}_2|_{W^1_1(\Gamma)} \\
&\le \alpha C_E C_A \left(1 + \beta C_A\right) \beta C_A |v|_{W^1_p(\Omega)} |r_1 - r_2|_{W^1_q(\Omega)}\,.
\end{aligned}$$

Invoking (4.16) along with $\theta_1 \ge \beta C_A/(1 + \beta C_A)$ we obtain

$$|\widetilde{r}_1 - \widetilde{r}_2|_{W^1_q(\Omega)} \le \left(1 - \theta_1\right) \beta C_A |r_1 - r_2|_{W^1_q(\Omega)} \le \frac{\beta C_A}{1 + \beta C_A} |r_1 - r_2|_{W^1_q(\Omega)}\,.$$

Therefore $T = T_2 \circ T_1 : \mathbb{V} \to \mathbb{V}$ is a contraction in $\mathbb{V}$. □

5.3. Second order sufficient condition. The final step is to prove the second order sufficient condition for the optimal control $\bar{u}$ found earlier, which in turn guarantees that $\bar{u}$ is locally unique. This hinges on an additional condition on $|v|_{W^1_p(\Omega)}$ depending on the parameter θ_3 given by

$$\begin{aligned}
\theta_3 = \frac{\theta_2^3}{\alpha^2} &\left(C_A\left(1 + \beta C_A\right)\left(1 + 2\beta C_A\right)\right)^{-1} \\
&\cdot \left(\alpha C_E \left(1 + \beta C_A\right)\Big(1 + \|\gamma_d\|_{L^2(\Gamma)}\Big) + \beta \lambda_y \left(\frac{1 - \theta_1}{\alpha C_E C_A} + \|y_d\|_{L^2(\Omega)}\right)\right)^{-1}.
\end{aligned} \tag{5.7}$$

THEOREM 5.6 (second order sufficient conditions). *If θ_1, θ_2 satisfy* (4.16), (4.19), *and in addition*

$$|v|_{W_p^1(\Omega)} \le \frac{\theta_3 \lambda_u}{2}, \tag{5.8}$$

then

$$\mathcal{J}''(\bar{u})h^2 \ge \frac{\lambda_u}{2} \|h\|^2_{L^2(\Gamma)} \qquad \forall h \in \mathcal{C}(\bar{u}). \tag{5.9}$$

Proof. Since G_v is twice Fréchet differentiable, according to Theorem 4.15, we can write the second order Fréchet derivative of $\mathcal{J}$ from (5.1) at $\bar{u}$ in the direction h^2 as

$$\mathcal{J}''(\bar{u})h^2 = \mathcal{J}_1''(G_v(\bar{u}))(G_v'(\bar{u})h)^2 + \mathcal{J}_1'(G_v(\bar{u}))G_v''(\bar{u})h^2 + \mathcal{J}_2''(\bar{u})h^2. \tag{5.10}$$

By recalling that $(\gamma_u, y_u) = G_v'(\bar{u})h$ solves (4.33) and $(\gamma_{uu}, y_{uu}) = G_v''(\bar{u})h^2$ solves (4.37), we can write

$$\mathcal{J}''(\bar{u})h^2 = \mathcal{J}_1''(G_v(\bar{u}))(\gamma_u, y_u)^2 + \mathcal{J}_1'(G_v(\bar{u}))(\gamma_{uu}, y_{uu}) + \mathcal{J}_2''(\bar{u})h^2,$$

where

$$\begin{aligned}
\mathcal{J}_1''(G_v(\bar{u}))(\gamma_u, y_u)^2 &= \|\gamma_u\|^2_{L^2(\Gamma)} + \lambda_y \|y_u\|^2_{L^2(\Omega)}, \\
\mathcal{J}_1'(G_v(\bar{u}))(\gamma_{uu}, y_{uu}) &= \int_\Gamma (\bar{\gamma} - \gamma_d)\gamma_{uu} + \lambda_y \int_\Omega (\bar{y} + v - y_d) y_{uu}, \\
\mathcal{J}_2''(\bar{u})h^2 &= \lambda_u \|h\|^2_{L^2(\Gamma)}.
\end{aligned}$$

This yields

$$\begin{aligned}
\mathcal{J}''(\bar{u})h^2 \ge{}& \lambda_u \|h\|^2_{L^2(\Gamma)} + \|\gamma_u\|^2_{L^2(\Gamma)} + \lambda_y \|y_u\|^2_{L^2(\Omega)} \\
&- \|\bar{\gamma} - \gamma_d\|_{L^2(\Gamma)} |\gamma_{uu}|_{W^1_\infty(\Gamma)} - \lambda_y \|\bar{y} + v - y_d\|_{L^2(\Omega)} |y_{uu}|_{W_p^1(\Omega)},
\end{aligned}$$

because of the Poincaré inequalities $\|z\|_{L^2(\Omega)} \le |z|_{W_2^1(\Omega)} \le |z|_{W_p^1(\Omega)}$ for the unit square Ω and $\|\zeta\|_{L^2(\Gamma)} \le |\zeta|_{W_2^1(\Gamma)} \le |\zeta|_{W_p^1(\Gamma)}$ for the unit interval Γ.

Estimates (4.39) and (4.40) in Theorem 4.15 imply

$$\begin{aligned}
\mathcal{J}''(\bar{u})h^2 \ge{}& \lambda_u \|h\|^2_{L^2(\Gamma)} - \frac{\alpha^2}{\theta_2^3} C_A (1 + \beta C_A)(1 + 2\beta C_A) |v|_{W_p^1(\Omega)} \\
&\cdot \Big(\alpha C_E (1 + \beta C_A) \|\bar{\gamma} - \gamma_d\|_{L^2(\Gamma)} + \beta \lambda_y \|\bar{y} + v - y_d\|_{L^2(\Omega)} \Big) \|h\|^2_{L^2(\Gamma)}.
\end{aligned}$$

Furthermore, from $(\bar{\gamma}, \bar{y}) \in \mathbb{B}_v$ we obtain

$$\|\bar{\gamma} - \gamma_d\|_{L^2(\Gamma)} \le \|\bar{\gamma}\|_{L^2(\Gamma)} + \|\gamma_d\|_{L^2(\Gamma)} \le 1 + \|\gamma_d\|_{L^2(\Omega)},$$

and

$$\|\bar{y} + v - y_d\|_{L^2(\Omega)} \le |\bar{y} + v|_{W_p^1(\Omega)} + \|y_d\|_{L^2(\Omega)}$$

with

$$|\bar{y}+v|_{W_p^1(\Omega)} \le (1+\beta C_A)|v|_{W_p^1(\Omega)} \le \frac{1-\theta_1}{\alpha C_E C_A}.$$

In view of the definition of θ_3 above, this in turn implies

$$\mathcal{J}''(\bar{u})h^2 \ge \left(\lambda_u - \frac{|v|_{W_p^1(\Omega)}}{\theta_3}\right)\|h\|_{L^2(\Omega)}^2 .$$

Therefore, the smallness condition (5.8) on v yields (5.9). □

Corollary 5.7 (quadratic growth). *Under the assumptions of Theorem 5.6 there exist $\theta > 0$ such that for all $h \in \mathcal{C}(\bar{u})$ with $\|h\|_{L^2(\Gamma)} \le \theta$ we have*

$$\mathcal{J}(\bar{u}+h) \ge \mathcal{J}(\bar{u}) + \frac{\lambda_u}{8}\|h\|_{L^2(\Gamma)}^2 . \tag{5.11}$$

Proof. We refer to [19, Theorem 4.23]. □

Corollary 5.7 implies that there exists a unique local minimum $\bar{u}$ solution to our OC-FBP. Moreover, (5.11) is equivalent to

$$\left\langle \mathcal{J}'(u) - \mathcal{J}'(\bar{u}), u-\bar{u}\right\rangle_{L^2(\Gamma)\times L^2(\Gamma)} \ge \frac{\lambda_u}{4}\|u-\bar{u}\|_{L^2(\Gamma)}^2 \quad \forall u \in \bar{u}+\mathcal{C}(\bar{u}). \tag{5.12}$$

Acknowledgments. This research was supported in part by NSF grants DMS-0807811 and DMS-1109325. We thank Michael Hintermüller for pointing out reference [23] and many fruitful comments on the manuscript. We also thank Abner Salgado for useful discussions throughout this work.

REFERENCES

[1] *Mini-workshop: Control of Free Boundaries*, Oberwolfach Rep., 4 (2007), pp. 447–486. Abstracts from the mini-workshop held February 11–17, 2007, Organized by C. M. Elliott, M. Hinze and V. Styles, Oberwolfach Reports. Vol. 4, no. 1.

[2] *New directions in simulation, control and analysis for interfaces and free boundaries*, Oberwolfach Rep., 7 (2010), pp. 253–324. Abstracts from the workshop held January 31–February 6, 2010, Organized by C. M. Elliott, Y. Giga, M. Hinze and V. Styles, Oberwolfach Reports. Vol. 7, no. 1.

[3] H. Antil, R.H.W. Hoppe, and C. Linsenmann, *Path-following primal-dual interior-point methods for shape optimization of stationary flow problems*, J. Numer. Math., 15 (2007), pp. 81–100.

[4] M. K. Bernauer and R. Herzog, *Optimal control of the classical two-phase Stefan problem in level set formulation*, SIAM J. Sci. Comput., 33 (2011), pp. 342–363.

[5] J. C. De Los Reyes and M. Hintermüller, *A duality based semismooth Newton framework for solving variational inequalities of the second kind*, Interfaces Free Bound., 13 (2011), pp. 437–462.

[6] K. Deckelnick, G. Dziuk, and C. M. Elliott, *Computation of geometric partial differential equations and mean curvature flow*, Acta Numerica, 14 (2005), pp. 139–232.

[7] M. C. Delfour and J.-P. Zolésio, *Shapes and geometries*, vol. 22 of Advances in Design and Control, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, second ed., 2011. Metrics, analysis, differential calculus, and optimization.

[8] A. Ern and J. L. Guermond, *Theory and practice of finite elements*, vol. 159 of Applied Mathematical Sciences, Springer-Verlag, New York, 2004.

[9] R. Glowinski, *Numerical methods for nonlinear variational problems*, Scientific Computation, Springer-Verlag, Berlin, 2008. Reprint of the 1984 original.

[10] M. Hinze and S. Ziegenbalg, *Optimal control of the free boundary in a two-phase Stefan problem*, J. Comput. Phys., 223 (2007), pp. 657–684.

[11] ———, *Optimal control of the free boundary in a two-phase Stefan problem with flow driven by convection*, ZAMM Z. Angew. Math. Mech., 87 (2007), pp. 430–448.

[12] C. T. Kelley, *Iterative methods for optimization*, vol. 18 of Frontiers in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.

[13] T. Krupenkin and J.A. Taylor, *Reverse electrowetting as a new approach to high-power energy harvesting*, Nat. Commun., 2 (2011), p. 448.

[14] S. Repke, N. Marheineke, and R. Pinnau, *Two adjoint-based optimization approaches for a free surface Stokes flow*, SIAM J. Appl. Math., 71 (2011), pp. 2168–2184.

[15] H. L. Royden, *Real analysis*, Macmillan Publishing Company, New York, third ed., 1988.

[16] P. Saavedra and L. R. Scott, *Variational formulation of a model free-boundary problem*, Math. Comp., 57 (1991), pp. 451–475.

[17] J. Sokołowski and J. P. Zolésio, *Introduction to shape optimization*, vol. 16 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, 1992. Shape sensitivity analysis.

[18] L. Tartar, *An introduction to Sobolev spaces and interpolation spaces*, vol. 3 of Lecture Notes of the Unione Matematica Italiana, Springer, Berlin, 2007.

[19] F. Tröltzsch, *Optimal control of partial differential equations*, vol. 112 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2010. Theory, methods and applications, Translated from the 2005 German original by Jürgen Sprekels.

[20] KG van der Zee, EH van Brummelen, I. Akkerman, and R. de Borst, *Goal-oriented error estimation and adaptivity for fluid-structure interaction using exact linearized adjoints*, Computer Meths. in Appl. Mech. and Eng., (2010).

[21] S.W. Walker, B. Shapiro, and R.H. Nochetto, *Electrowetting with contact line pinning: Computational modeling and comparisons with experiments*, Physics of Fluids, 21 (2009), p. 102103.

[22] S. W. Walker, A. Bonito, and R. H. Nochetto, *Mixed finite element method for electrowetting on dielectric with contact line pinning*, Interfaces Free Bound., 12 (2010), pp. 85–119.

[23] J. Zowe and S. Kurcyusz, *Regularity and stability for the mathematical programming problem in Banach spaces*, Appl. Math. Optim., 5 (1979), pp. 49–62.